

# Biodiversity and biotechnological potential of microorganisms from mangrove ecosystems: a review

Hrudayanath Thatoi · Bikash Chandra Behera ·  
Rashmi Ranjan Mishra · Sushil Kumar Dutta

Received: 20 December 2011 / Accepted: 23 February 2012 / Published online: 2 June 2012  
© Springer-Verlag and the University of Milan 2012

**Abstract** Mangrove forests occurring at the interface of terrestrial and marine ecosystems represent a rich biological diversity of plants, animals and microorganisms. Microbes, being an important component of the mangrove environment, not only play a very critical role in creating and maintaining this biosphere but also serve as a source of biotechnologically valuable and important products. By participating in various steps of decomposition and mineralization of leaf litter, microbes make an essential contribution to the productivity of the mangrove ecosystem. They are able to recycle nutrients, produce and consume gases that affect global climate, destroy pollutants, treat anthropogenic wastes and can also be used for biological control of plant and animal pests. Microorganisms from mangrove environments are a major source of antimicrobial agents and also produce a wide range of important medicinal compounds,

including enzymes, antitumor agents, insecticides, vitamins, immunosuppressants, and immune modulators. However, the phylogenetic and functional description of microbial diversity in mangrove ecosystems has not been addressed to the same extent as for other environments. Even though the mangrove ecosystem is very rich in microbial diversity, less than 5% of species have been described; in many cases neither their ecological role nor their application potential is known. Recently developed technologies in molecular biology and genetics offer great promise to explore the potential of microbial diversity. Hence, the present paper makes an attempt to review the microbial diversity in mangrove ecosystems and explore their potential applications in various fields such as agriculture, pharmaceutical, industrial, environmental and medical sciences.

**Keywords** Microbial diversity · Mangrove · Ecological role · Biotechnological potential

---

H. Thatoi (✉)  
Department of Biotechnology,  
College of Engineering and Technology,  
Biju Pattnaik University of Technology,  
Techno-Campus,  
Bhubaneswar 751003, Odisha, India  
e-mail: hn\_thatoi@rediffmail.com

B. C. Behera  
Department of Biotechnology,  
North Orissa University,  
Baripada 757003, Odisha, India

R. R. Mishra  
Department of Biotechnology,  
MITS School of Biotechnology,  
Bhubaneswar 751024, Odisha, India

S. K. Dutta  
Department of Zoology,  
North Orissa University,  
Baripada 757003, Odisha, India

## Introduction

Mangroves are unique inter-tidal ecosystems of the tropical and sub-tropical regions of the world that support genetically diverse groups of aquatic and terrestrial organisms. Nearly 60–70% of the world's tropical and subtropical coastlines are covered with mangroves, which are known to be highly productive ecosystems of immense ecological value. Despite being fragile and sparsely distributed, these ecosystems are highly productive all over the world (Thatoi and Biswal 2008). They protect and stabilize coastal zones, nourish and nurture the coastal water with nutrients. These ecosystems are characterized by periodic tidal flooding, which makes environmental factors such as salinity and nutrient availability highly variable resulting in unique and

specific characteristics. Apart from flora and fauna, microbial diversity characterizes one of the important communities of these ecosystems. Because of the abundance of carbon and other nutrient contents, the mangrove ecosystem harbors a large number of microbial communities, which can adapt to the moderate saline and fluctuating environmental conditions. These microbial communities play an important role in cycling of nutrients such as carbon, nitrogen, sulfur and phosphorous, and thus control the chemical environment of the mangrove ecosystem (Alongi et al. 1993). Microbial activity is also responsible for major nutrient transformations within mangrove ecosystems (Alongi et al. 1993; Holguin et al. 1999). In tropical mangroves, bacteria and fungi constitute 91% of the total microbial biomass, whereas algae and protozoa represent only 7% and 2%, respectively (Alongi 1988). Complex interactions of these microbes maintain the harmony of different biogeochemical processes and sustain the nutritional status and ecological balance. Free living bacteria, fungi and yeasts have been reported to play a significant role in the formation of detritus in mangrove ecosystems (Maria and Sridhar 2002). Various groups of bacteria are usually present in the ecosystem (Holguin et al. 2001), performing various activities like photosynthesis, nitrogen fixation, methanogenesis (Das et al. 2006). Several studies have shown the uniqueness of mangrove sediments with respect to their microbial composition (Gray and Herwig 1996; Urakawa et al. 1999). Studies on microbial diversity in the mangrove sediments are important in understanding the process of biogeochemical cycling and pollutant removal (Roy et al. 2002). The microbial diversity of mangrove ecosystems can also provide information on their ecological role and unique biotechnological potential in the field of agriculture, industry, medicine and pharmaceuticals (Lageiro et al. 2007). During the past decade, the development of molecular techniques using nucleic acids has led to many new findings in studies of microbial ecology (Amann et al. 1995). This approach avoids the limitations of traditional culturing techniques for assessing microbial diversity in natural environments. Notwithstanding the existing knowledge of microbes and microbial processes, we are still only scratching the surface of microbial diversity, which needs to be explored for the judicious and gainful utilization of this nature's treasure. The particular conditions of a mangrove, and adaptation of bacterial species to such conditions, represents an important source of biotechnological potential resources to be exploited (Sivaramakrishnan et al. 2006). Microorganisms from mangrove ecosystems contain useful enzymes, proteins, antibiotics and salt tolerant genes of much biotechnological significance. This review is an attempt to consolidate the latest studies on mangrove microbial biodiversity and to explore the immense biotechnological potential of mangrove microbial flora in various fields.

## Microbial diversity of mangrove ecosystems

Biodiversity can be defined as the variability of life at the genetic, species and ecosystem levels of organization. The mangrove ecosystem provides a unique environment harboring diverse groups of microorganisms. All microbial forms, such as bacteria, fungi, cyanobacteria, microalgae, macroalgae and fungus-like protists, have been reported in this ecosystem. The common bacterial groups of the mangroves are sulfate-reducing (*Desulfovibrio*, *Desulfotomaculum*, *Desulfosarcina*, *Desulfococcus* sp., etc.), N<sub>2</sub>-fixing (*Azospirillum*, *Azotobacter*, *Rhizobium*, *Clostridium*, *Klebsiella* sp., etc.), phosphate-solubilizing (*Bacillus*, *Paenibacillus*, *Xanthobacter*, *Vibrio proteolyticus*, *Enterobacter*, *Kluyvera*, *Chryseomonas* and *Pseudomonas* sp., etc.), photosynthetic anoxygenic (*Chloronema*, *Chromatium*, *Beggiatoa*, *Thiopedia*, *Leucothiobacteria* sp., etc.) and methanogenic (*Methanococcus methylutens* sp., etc.) bacteria. In addition, various groups of fungi, such as ligninolytic, cellulolytic, pectinolytic, amylolytic and proteolytic fungi as well as actinomycetes are present in mangrove ecosystems (Kathiresan and Bingham 2001). Among the algae, groups like *Chlorophyta*, *Chrysochyta*, *Phaeophyta*, *Rhodophyta* and *Cyanophyta* dominate the mangrove ecosystem (Sen and Naskar 2003).

## Algae

In mangroves, algal species occur abundantly in planktonic, benthic and periphytic forms. The roots (pneumatophores) of mangrove trees provide a favorable habitat for 50% of the total algal species, the hard substrates for 30% and the soft mud for 20% of the species (Kathiresan and Qasim 2005). Certain algae are associated frequently with mangroves and are considered as characteristic of the ecology. Algal species belonging to the genera *Bostrychia*, *Cologlossa* and *Catenella* are commonly present on the roots and trunks of the mangroves. Species of the genera *Rhizoclonium*, *Enteromorpha* and *Cladophora* normally exist in the sediment along with the cyanobacteria (*Lyngbya* and *Anacystis*), benthic diatoms and the sulfate-reducing bacteria (Kathiresan and Qasim 2005). This microbial community forms a 'bio-film' that activates the attachment of algae to the mangrove trees. The factors governing the occurrence of these algae are mainly those of a microclimate prevailing in the swamps. Most of these algae are small filament forms that are fairly resistant to desiccation as well as to high salinity.

Algae are widespread in mangrove habitats all over the world. Cordeiro-Marino et al. (1992), recorded 150 algal taxa from New World mangroves, with highest diversity in the red algal groups with 78 species, and lowest in the brown algae with less than 15 species. Maximum algal diversity with 109 species and a high degree of endemism (about 70%) of red

algae was reported along the Caribbean coast (Kathiresan and Qasim 2005). In the Indian context, about 558 species of algae from seven families are known to occur (Kathiresan and Qasim 2005). The east coast is represented by 264 species, whereas the west coast has 326 species (Kathiresan and Qasim 2005). Only 71 species have been recorded so far from the Andaman and Nicobar Islands (Kathiresan and Qasim 2005). From the Sundarbans mangroves of West Bengal, 150 different species of algal flora were identified, belonging to different groups viz., *Chlorophyta* (39 species), *Chrysophyta* (44 species), *Phaeophyta* (2 species), *Rhodophyta* (15 species) and *Cyanophyta* (50 species) (Sen and Naskar 2003). Algal species such as *Gloeocapsa* sp., *Chlorella* sp., *Ulva* sp., *Anabaena* sp., *Spirogyra* sp., *Oscillatoria* sp., *Phormidium* sp. were identified from mangroves of Bhitarkanika, Orissa (Mishra 2010).

## Fungi

Mangrove forests are biodiversity ‘hotspots’ for marine fungi (Shearer et al. 2007). Although the mangrove trunks and aerating roots are submerged, permanently or intermittently, the upper parts of the roots and trunks are rarely or never wetted by the salt water. Thus, terrestrial fungi and lichens occupy the upper part of the trees and marine species occupy the lower part; at the interface there is an overlap between marine and terrestrial fungi (Sarma and Hyde 2001). Since they were first reported from mangrove roots in Australia by Cribb and Cribb (1995), there has been considerable increase in information on mangrove-associated fungi. The latest estimate of marine fungal species in the world is 1,500, which excludes lichens and many fungi that are newly isolated or inadequately described (Hyde et al. 1998). Out of large number of estimated fungal species, Hyde listed only 120 species of fungi from 29 mangrove forests around the world (Hyde 1990). These included 87 *Ascomycetes*, 31 *Deuteromycetes* and 2 *Basidiomycetes*. There are some 169 fungal species from Malaysia (Alias et al. 1995), 44 fungi associated with standing senescent *Acanthus ilicifolius* from Mai Po mangrove, Hong Kong (Sadaba et al. 1995), 76 species from Pearl River Estuary, China (Vrijmoed et al. 1991), 91 fungi from Egyptian Red Sea (Abdel-Wahab 2005) and 112 species reported from the Bahamas islands (Jones and Abdel-Wahab 2005). In the Indian context, about 48 fungal species were reported to occur in decomposing *Rhizophora* debris in Pichavaram, South India (Ravikumar and Vittal 1996). Seven species of fungi that exist on mangrove leaf surfaces in the Sundarbans of West Bengal have been reported by Pal and Purkayastha (1992). From a Mangalvan mangrove ecosystem, 31 fungal isolates were recorded from soil, and 27 species from decaying mangroves; 7 species from floating plants were reported, with dominance of *Aspergillus* followed by *Penicillium*, *Fusarium* and *Trichoderma*. Sarma and Hyde (2001)

reported 73 species of fungi from the Krishna estuaries of India. Similarly, 31 fungal species have been studied from sediment and 27 species from decaying leaves, stems, roots and pneumatophores of an estuarine mangrove ecosystem in Cochin (Prabhakaran et al. 1990). Raghukumar et al. (1995) studied the colonization of thraustochytrids on leaf litter of *Rhizophora apicula* at Caorao mangrove, Goa. Higher groups of fungi have been reported from the mangrove woods on the Maharashtra coast, with 41 species of *Ascomycetes*, 2 *Basidiomycetes*, and 12 *Deuteromycetes* with predominance of *Massarina velatospora* (Borse 1988). In Bhitarkanika, Orissa (India), Gupta et al. (2009a, b) reported the population status of fungi associated with the phyllosphere of different mangrove plants viz., *Avicennia*, *Aegiceras*, *Bruguiera*, *Ceriops*, *Excoecaria*, *Heritiera*, *Kandelia*, *Rhizophora* and *Sonneratia*. A total of 33 fungi from Godavari and 67 fungi from the Krishna estuary, India, was reported by Venkateswara Sarma et al. (2001). Relatively few fungi have been reported as pathogens of mangrove plants compared to the number of saprophytic fungi identified on decaying mangrove wood and leaves (Hyde et al. 1998). The intertidal fungus *Cytospora rhizophorae* is thought to be parasitic on *Rhizophora* spp. prop root (Kohlmeyer and Kohlmeyer 1979). *Phomopsis mangrovei*, which is probably pathogenic, was described from dying prop roots of *Rhizophora apiculata* in Thailand (Hyde 1996). An intertidal *Phytophthora* species was described to cause terminal dieback of *Avicennia marina* (Pegg et al. 1980; Gadek 1998) and a *Phytophthora* species was also found to be pathogenic on *Avicennia marina* var. *resinifera* in New Zealand (Maxwell 1968). *Halophytophthora* species were also thought to be responsible for diseased mangrove forests over vast areas in Sydney (Garrettson-Cornell and Simpson 1984).

Endophytic fungi were found in large numbers in mangrove environments. More than 200 species of endophytic fungi—mainly *Alternaria*, *Aspergillus*, *Cladosporium*, *Colletotrichum*, *Fusarium*, *Paecilomyces*, *Penicillium*, *Pestalotiopsis*, *Phoma*, *Phomopsis*, *Phyllosticta* and *Trichoderma*—were isolated and identified from mangroves (Liu et al. 2007). Chinese researchers surveyed and reported the occurrence of several arbuscular mycorrhizal fungi (AMF) on root system mangroves plants in QinZhou Bay, Guangxi, China (Wang et al. 2003). In India, 25 endophytic fungi comprised of 3 *ascomycetes*, 20 *mitosporic* fungi and 2 *sterile* fungi were recovered from 2 halophytes (*Acanthus ilicifolius* and *Acrostichum aureum*) of a west coast mangrove habitat. Endophytic fungi were also isolated from leaves of *Rhizophora apiculata* and *Rhizophora mucronata*, two typical mangrove plants grown in the Pichavaram mangrove of Tamil Nadu, Southern India (Suryanarayanan et al. 1998). Vazquez et al. (2000) first isolated a marine phosphate-solubilizing fungus, *Aspergillus niger*, together with several phosphate-solubilizing bacterial strains, from the rhizosphere of black mangrove *Avicennia germinans*,

and proposed the production of organic acids by these mangrove rhizosphere microorganisms as a possible mechanism involved in the solubilization of insoluble calcium phosphate. Arbuscular mycorrhizal (AM) colonizing mangrove and mangrove associates from river estuary of Ganges have been reported by Sengupta and Chaudhuri (2002).

### Actinomycetes

Actinomycetes have been looked upon as potential sources of bioactive compounds and are the richest sources of secondary metabolites. The mangrove ecosystem is a largely unexplored source of actinomycetes with the potential to produce biologically active secondary metabolites (Hong et al. 2009). Several reports from different geographical locations around the world have described the occurrences of actinomycetes in different mangrove habitats. Eccleston et al. (2008) reported the occurrence of actinomycetes belongs to genus *Micromonospora* from the Sunshine Coast in Australia. A rifamycin-producing *Micromonospora* from mangroves of the South China Sea was reported by Huang et al. (2008) and Xie et al. (2006). Several genera of actinomycetes, such as *Actinomadura*, *Microbispora*, *Nonomuraea*, *Actinoplanes*, *Micromonospora*, *Verrucosipora*, *Arthrobacter*, *Isoptericola*, *Micrococcus*, *Microbacterium*, *Nocardia*, *Rhodococcus* and *Streptomyces* were reported from mangrove soils and plants in China (Hong et al. 2009). Similarly, genera like *Brevibacterium*, *Dermabacter*, *Kocuria*, *Kytococcus*, *Microbacterium*, *Nesterenkonia* and *Rothia* were reported from a mangrove sediment in Brazil (Dias et al. 2009). Ara et al. (2007) reported novel actinomycetes (*Nonomuraea maheshkhaliensis*) from a mangrove rhizosphere mud in the southern area of Bangladesh.

In the Indian context, Sivakumar (2001) reported 23 actinomycete species from Pichavaram mangrove, with most species identified belonging to the genus *Streptomyces*. Laksmanaperumalsamy et al. (1978) isolated 518 *Streptomyces* strains from mangrove environments of Porto Novo. As many as 107 different actinomycetes were isolated from marine sediments off the Konkan coast of Maharashtra by Gulve and Deshmukh (2011), and 17 actinomycetes isolates were identified from the Karangkadu mangrove forest of Tamil Nadu, India by Ravikumar et al. (2011b). Sahu et al. (2005) reported several *Streptomyces*, viz., *Streptomyces alboniger*, *S. violaceus*, *S. moderatus* and *S. aureofasciculus* from the Vellar estuary on the south east coast of India. Similarly, Ravikumar et al. (2011a, 2011b) reported the biodiversity of actinomycetes from sediments of the Manakkudi mangrove ecosystem of the southwest coast of India. Sivakumar et al. (2005) reported the occurrence of *Streptomyces albidoflavus*, which has antitumor properties, from the Pichavaram mangrove. A number of *Streptomyces*, such as *S. albidoflavus*, *S.*

*atroolivaceus*, *S. auranticus*, *S. canus*, *S. chromofuscus*, *S. exfoliates*, *S. griseoluteus*, *S. helstedii*, *S. lavenduale*, *S. longisporoflavus*, *S. luridus*, *S. lydicus*, *S. nogalator*, *S. pactum*, *S. prasinosporus*, *S. purpureus*, *S. tubercidus*, *S. versoviensis*, *S. viridochromogenes* and *S. xanthochromogenes* have been reported by Gupta et al. (2009a, b) from different plant species of the Bhitarkanika mangrove forest of Orissa, India. Phosphate-solubilizing actinomycetes such as *Streptomyces galbus* in the Vellar Estuary of the Parangipettai estuarine environment on the south east coast of India have been reported by Sahu et al. (2007), and five phosphate-solubilizing *Streptomyces* spp. from the mangrove ecosystem of Bhitarkanika, Orissa, India were reported by Gupta et al. (2010). Distribution of actinomycetes in the Sundarbans mangrove of West Bengal, India has been reported by Mitra et al. (2008). This relatively large distribution of actinomycetes species throughout the world's mangrove ecosystems appears to reason that mangrove forests represent a treasure trove of actinomycetes.

### Bacteria

Next to trees, bacterial flora dominates the biomass and productivity of mangrove forests (Kathiresan and Qasim 2005). Among microbes, bacterial populations in mangroves are many fold greater than those of fungi (Kathiresan and Qasim 2005). Microbially generated detritus in mangrove ecosystems acts as the major substrate for bacterial growth in mangroves (Bano et al. 1997). The bacteria may act as primary decomposers that utilize dissolved organic substances at low concentration and assimilate dissolved inorganic substances like nitrate and phosphate. Bacteria with different ecological roles have been reported from mangrove ecosystems.

#### N<sub>2</sub> fixing bacteria

Nitrogen fixation is a process of conversion of gaseous forms of nitrogen (N<sub>2</sub>) into combined forms, i.e., ammonia or organic nitrogen by some bacteria and cyanobacteria. Free-living as well as symbiotic microbes known as diazotrophs fix N<sub>2</sub> into proteins. Nitrogen-fixing (diazotrophic) microorganisms can colonize both terrestrial and marine environments. In mangrove ecosystems, high rates of nitrogen fixation have been associated with dead and decomposing leaves (Mann and Steinke 1992), pneumatophores (Hicks and Silvester 1985; Toledo et al. 1995) and the rhizosphere soil (Holguin et al. 1992). N<sub>2</sub> fixation in mangrove sediments is likely to be limited by insufficient energy sources. The low rates of N<sub>2</sub> fixation by heterotrophic bacteria detected in marine water are probably due to lack of energy sources. Nitrogen fixation by heterotrophic bacteria can be regulated

by specific environmental factors such as oxygen, combined nitrogen and the availability of carbon source to support energy requirements. Energy for  $N_2$  fixation can also be derived from leaves and roots decomposed by non-diazotrophic microflora that colonize dead mangrove leaves (Zuberr and Silver 1978). Nitrogen-fixing bacteria such as members of the genera *Azospirillum*, *Azotobacter*, *Rhizobium*, *Clostridium* and *Klebsiella* were isolated from the sediments, rhizosphere and root surfaces of various mangrove species. Nitrogen-fixing bacteria, *Azotobacter* species are repeatedly isolated from sediments of Pichavaram mangroves, and they were more in the mangrove habitats than in the backwaters and estuarine systems (Lakshmanaperumalsamy 1987). Several strains of diazotrophic bacteria such as *Vibrio campbellii*, *Listonella anguillarum*, *Vibrio aestuarianus* and *Phyllobacterium* sp. were isolated from the rhizosphere of mangroves in Mexico (Holguin et al. 1992).  $N_2$ -fixing bacteria such as *Azotobacter* sp. which can be used as biofertilizers, are abundant in mangrove habitats of Pichavaram (Ravikumar 1995). Two halotolerant  $N_2$ -fixing *Rhizobium* strains were isolated from root nodules of *Derris scandens* and *Sesbania* species growing in the mangrove swamps of Sundarbans (Sengupta and Choudhury 1990). Nitrogen-fixing cyanobacteria such as *Aphanocapsa* sp., *Nodularia* sp., and *Trichodesmium* sp., were isolated from Pichavaram mangroves (Ramachandran and Venugopalan 1987).  $N_2$ -fixing bacteria are efficient at using a variety of mangrove substrates despite differences in carbon content and phenol concentrations (Pelegri and Twilley 1998). However, their abundance may be dependent on physical conditions and mangrove community composition. Both symbiotic and asymbiotic  $N_2$ -fixing bacteria play a vital role in the nitrogen enrichment of mangrove ecosystems (Holguin et al. 2001). One may conclude from the available information that  $N_2$  fixation is a major bacterial activity in mangrove ecosystems, second only to carbon decomposition of detritus by sulfate-reducing bacteria.

#### Phosphate-solubilizing bacteria

Muddy mangrove soils have a strong capacity to absorb nitrates and phosphates carried by the tides (Hesse 1962). Most of the inorganic phosphate present in the sediment is bound to calcium, iron and aluminium ions as insoluble phosphates (Alongi et al. 1992). Fungi and inorganic phosphate-solubilizing bacteria present in the mangrove rhizosphere participate in releasing soluble phosphate into pore water (Vazquez et al. 2000). Certain bacteria exhibit high phosphatase activity, capable of solubilizing phosphate (Sundararaj et al. 1974). Phosphorous is one of the major plant nutrients, second only to nitrogen (Vassileva et al. 1998), so phosphate-solubilizing microorganisms (PSMs) play an important role in supplementing phosphorus to plants and allowing the sustainable use of phosphate

fertilizers (Gyaneshwar et al. 1998). In an arid mangrove ecosystem in Mexico, nine strains of phosphate-solubilizing bacteria, such as *Bacillus amyloliquefaciens*, *B. atrophaeus*, *Paenibacillus macerans*, *Xanthobacter agilis*, *Vibrio proteolyticus*, *Enterobacter aerogenes*, *E. taylorae*, *E. asburiae* and *Kluyvera cryocrescens* were isolated from black mangrove (*Avicennia germinans*) roots. A further three strains, viz. *B. licheniformis*, *Chryseomonas luteola* and *Pseudomonas stutzeri* were isolated from white mangrove (*Laguncularia racemosa*) roots (Vazquez et al. 2000). Very little information is available about phosphate-solubilizing bacterial diversity and their activity in Indian mangroves. However, some studies related to phosphate-solubilizing bacterial activity have been done in the Vellar estuary at Parangipettai on the south eastern coast of India (Kathiresan and Selvam 2006), the Bhitarkanika mangrove environment of Orissa (Gupta et al. 2007; Mishra 2010), the Sundarban mangroves of West Bengal (Ramanathan et al. 2008) and the Great Nicobar mangroves of India (Kothamasi et al. 2006). A preliminary isolation and screening of phosphate-solubilizing bacteria from mangrove soil from Bhitarkanika, on the Orissa coast, by Gupta et al. (2007) revealed the presence of 33 soil bacteria showing phosphate-solubilizing capacity. Ramanathan et al. (2008) quantified phosphorus-solubilizing bacteria along with cellulose-degrading and  $N_2$ -fixing bacteria from Sundarban mangroves of India. Kothamasi et al. (2006) reported two strains of phosphate-solubilizing *Pseudomonas aeruginosa* (designated GM01 and GM02) in mangrove soils of Great Nicobar. Nine phosphate-solubilizing bacteria have been isolated and characterized phenotypically from mangrove soil of Bhitarkanika (Mishra 2010). Genera of phosphate-solubilizing bacteria, like *Pseudomonas*, *Bacillus*, *Corynebacterium*, *Vibrio*, *Micrococcus* and *Alcaligenes*, were studied by Venkateswaran and Natarajan (1983) in mangrove biotopes in Porto Novo, Chennai water and sediment. Endophytic phosphate-solubilizing bacteria were isolated from leaf samples of mangrove plants of Pichavaram, Tamil Nadu by Gayathri et al. (2010).

#### Sulfur-oxidizing bacteria

Bacteria also play a major role in the chemical and biological redox reactions that create the sulfur cycle. Sulfur and sulfide-oxidizing bacteria generate sulfate, which is used by sulfate-reducing bacteria (SRB) as an alternative electron acceptor in anaerobic respiration to create hydrogen sulfide. SRB cycle hydrogen sulfide through the atmosphere for use by anaerobic photosynthetic bacteria and sulfur-oxidizing bacteria, while returning carbon dioxide to the atmosphere (Holmer and Storkholm 2001). Mangrove sediments are mainly anaerobic with an overlying thin aerobic sediment layer. Degradation of organic matter in the aerobic zone occurs principally through

aerobic respiration, whereas in the anaerobic layer decomposition occurs mainly through sulfate-reduction (Sherman et al. 1998). Sulfur-oxidizing bacteria play an important role in the detoxification of sulfide in sediments. Symbiotic sulfur-oxidizers, e.g., those within members of the bivalve family Lucinacea, are commonly found in muddy mangrove areas (Liang et al. 2006). Sulfate reduction accounts for almost 100% of the total emission of CO<sub>2</sub> from the sediment (Kristensen et al. 1991). Some sulfur-oxidizing bacteria like Gammaproteobacteria, e.g., *Chromatiales*, and Deltaproteobacteria, e.g., *Desulfobacterales*, were reported from oil-contaminated soil of Brazilian pristine mangrove sediment (Santos et al. 2011; Holguin et al. 2001). The bacterial diversity present in sediments of a well-preserved mangrove in Ilha do Cardoso, located in the extreme south of the São Paulo State coastline, Brazil, was assessed using culture independent molecular approaches [denaturing gradient gel electrophoresis (DGGE)]. The data revealed a bacterial community dominated by Alphaproteobacteria (40.36% of clones), Gammaproteobacteria (19.28% of clones) and Acidobacteria (27.71% of clones) besides minor components of Firmicutes, Actinobacteria and Bacteroidetes (Dias et al. 2010). Some free-living and symbiotic sulfur oxidizing bacteria were reported from the Futian mangrove swamp of China (Liang et al. 2006). In Florida, SRB were the most numerous bacterial group in the rhizosphere of *Rhizophora mangle* and *A. germinans* mangroves, reaching a population density of 10<sup>6</sup> cfu g<sup>-1</sup> fresh weight (Zuberr and Silver 1978). In Goa's mangrove (India), 10<sup>3</sup> cfu g<sup>-1</sup> dry sediment of SRB, mostly spore-forming species, were associated with mangroves (Saxena et al. 1988). Further, In Goa's mangroves, eight species of SRB such as *Desulfovibrio desulfuricans*, *Desulfovibrio desulfuricans aestuarii*, *Desulfovibrio salexigens*, *Desulfovibrio sapovorans*, *Desulfotomaculum orientis*, *Desulfotomaculum acetoxidans*, *Desulfosarcina variabilis* and *Desulfococcus multivorans* were isolated and tentatively classified within four different genera (Loka Bharathi et al. 1991). In mangrove sediments, availability of iron and phosphorus may also depend on the activity of SRB (Holguin et al. 2001). It appears that, as the main decomposers of organic matter in anaerobic sediments, SRB play a major role in the mineralization of organic sulfur and production of the soluble iron and phosphorus used by organisms in mangrove ecosystems.

#### Cellulose-degrading bacteria

Cellulose is the primary product of photosynthesis in terrestrial environments and the most abundant renewable biorenewable product in the biosphere (Zhang and Lynd 2004). Cellulose biodegradation by cellulases and cellulosomes, produced by numerous microorganisms, represents the major carbon flow from fixed carbon sinks to atmospheric CO<sub>2</sub>, and is very important in several agricultural and waste

treatment processes (Haight 2005). In anaerobic environments, which are rich in decaying plant material, decomposition of cellulose is brought about by complex communities of interacting microorganisms (Odum and Heald 1972). As the substrate, i.e., cellulose, is insoluble, bacterial and fungal degradation occurs exocellularly to degrade cellulose into carbon and energy sources that are required by other microorganisms present in the mangrove environment. Several marine bacterial species such as *Rhodospirillum rubrum*, *Cellulomonas fimi*, *Clostridium stercorarium*, *Bacillus polymyxa*, *Pyrococcus furiosus*, *Acidothermus cellulolyticus*, *Saccharophagus degradans* have been reported as degrading cellulose (Taylor et al. 2006). Five promising cellulose-producing bacteria such as *Bacillus cereus*, *Bacillus licheniformis*, and *Bacillus pumilus* and *Bacillus* sp., have been reported from mangroves in the Philippines (Tabao and Moasalud 2010). Sediment associated with dense Sunderban mangroves showed highest counts of cellulose-degrading bacteria in comparison to other bacterial diversity (Ramanathan et al. 2008). Compared to other bacterial groups, much less information is available on the diversity of cellulose-degrading bacteria from mangrove ecosystems, which may be due to the lack of suitable technologies for their isolation and identification. A detailed study is required to assess the diversity of cellulose-degrading bacteria from various mangrove ecosystems.

#### Photosynthetic anoxygenic bacteria

Two main types of photosynthetic bacteria are seen in mangrove ecosystems, viz., purple sulfur bacteria (PSB; family Chromatiaceae), and purple non-sulfur bacteria (PNB; family Rhodospirillaceae, strains belonging to *Rhodospseudomonas* spp.). These bacteria are capable of using light to grow, fix nitrogen and release hydrogen gas in this environment. PSB range in color from pink to purple and contain bacteriochlorophyll *a* as their major pigment. These phototrophic anaerobes require sulphide, which they oxidise to sulphate for growth. Carbon dioxide is the usual source of cell carbon, but they also utilise various organic acids as carbon sources and are usually distributed widely in sulfide-rich reducing environments such as mangroves (Vethanayagam 1991). PNB range in color from brown to red and also contain bacteriochlorophyll *a* as their major pigment. They have the ability to utilize a remarkably wide spectrum of reducing carbon compounds, like malate or succinate, as electron donors as well as carbon sources for growth. Sulfur-rich mangrove ecosystems, with their mainly anaerobic soil environment, provide favorable conditions for the proliferation of these bacteria. The predominant bacteria belonging to this group in the mangrove ecosystem of Cochin (India) were identified as members of the genera *Chloronema*, *Chromatium*, *Beggiatoa*,

*Thiopedia* and *Leucothiobacteria* (Vethanayagam and Krishnamurthy 1995). In mangrove on the coast of Red Sea in Egypt, 225 isolates of PNB, belonging to 10 species of 4 different genera, were identified. Nine of the ten species inhabited the rhizosphere and root surface of the trees. The most common bacteria (*Rhodobacter* and *Rhodospseudomonas*) were detected in 73% and 80% of the sample, respectively (Shoreit et al. 1994). Some of the anoxygenic photosynthetic bacteria were also diazotrophic. Although there is as yet no published evidence, one can hypothesize that photosynthetic anoxygenic bacteria—the predominant photosynthetic organisms in anaerobic environments—contribute substantially to ecosystem productivity (Saho and Dhal 2009).

#### Methanogenic bacteria


An important characteristic of mangrove sediments is the absence of oxygen at a few millimeters below the surface (Lyimo et al. 2002). This lack of oxygen, coupled with the abundance of organic matter, creates an optimal environment for several groups of anaerobic organisms, such as SRB and methanogens (Dar et al. 2008). Since the presence of these groups in most coastal sediments is selected by the redox potential (Dar et al. 2008), these groups are expected to be found in discrete niches. However, these organisms are known to share similar niches in organic matter (OM)-rich environments like mangroves (Oremland et al. 1982). Niche superimposition between SRBs and methanogens is restricted to certain substrates, such as hydrogen and acetate (Oremland et al. 1982). Simple substrates (e.g., methanol, mono-di-trimethylamine) are important for methanogens (Lyimo et al. 2002), but not for SRB, which are capable of degrading more complex substrates, such as long-chain and aromatic hydrocarbons (Muyzer and Stams 2008). The presence of SRB limits the proliferation of these bacteria (Ramamurthy et al. 1990). There are several reports of methanogenic bacteria occurring in mangrove ecosystems, such as a strain of methanogenic bacterium, *Methanococcoides methylutens* (Mobanraju et al. 1997) and four strains of unidentified thermotolerant methanogenic bacteria isolated from the sediment of a mangrove forest (Marty 1985). A methanogenic bacterium, *Methanococcoides methylutens*, was isolated and characterized from the sediment of the mangrove environment of Pichavaram, southeast India (Mobanraju et al. 1997). Lyimo et al. (2008) also reported the methanogenic bacteria *Methanococcoides methylutens* and *Methanosarcina semesiae* from sediment samples of Tanzanian mangrove, and Taketani et al. (2010) reported the occurrence of methanogenic bacteria such as *Methanopyrus kandleri* and *Methanothermococcus thermolithotrophicus* from a pristine tropical mangrove soil of Brazil.

#### Ecological role of microorganisms in mangrove environment

Mangrove ecosystems provide large quantities of organic matter to adjacent coastal waters in the form of detritus, which serves as a nutrient source and is the base of an extensive food web. The diverse microbial communities living in mangrove ecosystems continuously transform nutrients from dead mangrove vegetation into sources of nitrogen, phosphorus and other nutrients that can be used by mangrove plants (Fig. 1). In turn, plant root exudates serve as a food source for the microorganisms living in the ecosystem. The degradation of mangrove vegetative material that produces detritus can be defined as organic matter in the active process of decomposition. It is rich in energy and contains a large active microbial population—both attached and free living (Odum and Heald 1975a). In addition to bacteria and fungi, other organisms may also colonize the vegetative material and contribute to detritus formation (D’Croz et al. 1989). Microscopic examination of decomposing mangrove leaves reveals a complex community composed of bacteria, fungi, protozoa and microalgae (Odum and Heald 1975b).

Bacteria are responsible for most of the carbon flux in tropical mangrove sediments. The element carbon, which forms the basis of all organic matter, undergoes a constant cycle in nature by various heterotrophic bacteria. The latter process most of the energy flow and nutrients and act as a carbon sink. Roughly 30–50% of the organic matter in mangrove leaves are leachable, containing water soluble compounds such as tannins and sugars (Cundell et al. 1979). The remaining fraction of organic matter consists of structural polymers such as lignocelluloses, which are used directly by the bacteria present in the soil and sediment of the mangrove. Nitrogen—a constituent of protein—is cycled in aquatic environments by several bacteria. The nitrate present in sediments, and derived from the degradation of nitrogenous organic compounds, is probably converted to ammonium ions by bacteria and is then assimilated by plants and bacteria. This process conserves the nitrogen within the ecosystem (Rivera-Monroy et al. 1995a, b). However, high rates of denitrification have been found in mangrove ecosystems into which wastewater are discharged, suggesting a correlation between denitrification rates and nitrate concentration (Rivera-Monroy et al. 1995b). Fixation of molecular nitrogen is carried out intracellularly by various bacteria, e.g., *Azotobacter*, *Clostridium*, etc. Some mangrove ecosystems are nutrient deficient especially in nitrogen and phosphorus (Sengupta and Choudhury 1991). Usually, phosphates precipitate because of the abundance of cations in the interstitial water of mangrove sediments, making phosphorus largely unavailable to plants. As potential suppliers of soluble forms of phosphorus, phosphate-

**Fig. 1** Conceptual model of ecological role of microbes in a mangrove ecosystem



solubilizing bacteria would have a great advantage for mangrove plants. Microorganisms also play an important part in sulfur transformations in mangrove ecosystems. Mangrove sediments are mainly anaerobic, with a thin overlying aerobic sediment layer. Degradation of organic matter in the aerobic zone occurs principally through aerobic respiration, whereas in the anaerobic layer decomposition occurs mainly through sulfate-reduction (Nedwell et al. 1994; Sherman et al. 1998). Previous investigations also suggest that sulfate-reduction may be an important pathway of organic matter mineralization in the organic-rich deposits typical of mangrove forests (Alongi et al. 1998). When sulfate is reduced by SRB, soluble sulfur compounds such as  $H_2S$  and  $HS$  are produced. These soluble sulfur compounds react with iron, reducing Fe (III) to Fe (II) and yielding pyrite ( $FeS_2$ ). Reduction of Fe (III) to Fe (II) releases soluble phosphate. It appears that SRB, as the main decomposers of organic matter in anaerobic sediments, play a major role in the mineralization of organic sulfur and in the production of soluble iron and phosphorus used by organisms in mangrove ecosystems. SRB can also contribute to the well-being of the ecosystem by fixing  $N_2$  (Holguin et al. 2001).

Fungi are decomposers of dead organic substrates associated with the decay of mangrove leaves (Fig. 1) (Nakagiri et al. 1996). Their importance lies in their ability to aggressively degrade lignocelluloses. They may be important in the degradation of dead animals and animal parts. Marine fungi are

important pathogens of plants and animals and also form symbiotic relationships with other organisms (Hyde et al. 1998). Mangrove fungi decompose vegetative material and thereby allow secondary colonization by bacteria and yeasts that further decompose the organic matter (Matondkar et al. 1981). In an Indian mangrove, the first colonizers of fallen mangrove leaves were fungi and thraustochytrids (fungi-like unicellular protists). Degradation of fallen mangrove vegetation starts immediately after its colonization by fungi and bacteria residing in the sediment and may last for 2–6 months, or more for degradation of wood (Steinke et al. 1990).

Phytoplankton and benthic micro-algal communities make an important contribution to the functioning of the mangrove environment. Due to the tidal influence and mixing of fresh and marine water in the mangrove ecosystem, several species of green-, and blue-green algae and seaweeds enter mangrove waters. Certain algae, which are colonised frequently and submerged in the surface area of sediments, roots, aerial roots, branches and trunks of mangrove, constitute one of the commercially most important marine resources, having the unique characteristic of fixing atmospheric carbon and nitrogen (Nedumaran et al. 2008). Algae and cyanobacteria also act as primary producers in the food chain of mangrove ecosystems. They harness solar energy and transform inorganic substances into food for other organisms. The consumption of food (phytoplankton) by the benthic organism, with their rapid assimilation of local carbon sources highlights the

importance of mangrove algal species in influencing carbon cycling and increasing the productivity in estuaries. Algal flora probably plays an important role in stabilizing and improving the physical properties of the soil by aggregating particles and adding organic matter to the ecosystem.

Besides bacteria, algae and fungi, actinomycetes are also reported to contribute to the breakdown and recycling of organic compounds. The production of several enzymes, such as cellulolytic, proteolytic, amylolytic, lipolytic, chitinolytic, phosphate-solubilizing activities, has been reported from marine actinomycetes (Sivakumar et al. 2007), which play a role in biodegradation of organic matter, thereby enhancing the productivity of the marine environment. Along with the phosphatase enzyme, actinomycetes play a major role in solubilizing phosphate in estuarine ecosystems and increasing the soluble phosphate concentration, thereby enhancing plant growth and increasing productivity (Sahu et al. 2007). Actinobacteria also play an important role by supplying food for other organisms by producing secondary metabolites that enhance the growth of juvenile fish, shrimp and prawn in the ecosystem (Sivakumar et al. 2007).

### Biotechnological potential of mangrove microorganisms

Microbial diversity is the key to human survival and economic wellbeing and provides a huge reservoir of resources that we can utilize for our benefit. Mangrove microorganisms have proven to be an important source of food, feed, medicine, enzymes and antimicrobial substances (Lin et al. 2001; Maria et al. 2005). Both halotolerant and halophilic bacteria and other microbes from mangrove ecosystem have large numbers of industrial applications in terms of their unique enzymes (Sabu 2003) that are capable of producing biosurfactants (Yakimov et al. 1999), bioplastics (Steinbüchel and Fuchtenbusch 1998), compatible solutes (Margesin and Schinner 2001), natural bioproducts and other commercially important products. Filamentous fungi—the principal commercial sources of xylanolytic enzymes—have many industrial uses, such as in paper manufacturing, animal feed, bread making, juice preparation, the wine industry and in xylitol production (Polizeli et al. 2005). Marine algae are the only source of industrially important phycocolloids like agar, carrageenan and alginate (Shanmugam and Mody 2000). They are also reported to have blood anticoagulant, anti-tumor, anti-mutagenic, anti-complementary, immunomodulating, hypoglycemic, antiviral, hypolipidemic and anti-inflammatory activities. Actinomycetes isolated from mangrove habitats are a potentially rich source of anti-infection and anti-tumor compounds and of agents for treating neurodegenerative diseases and diabetes

(Hong et al. 2009). Besides the above, mangrove microorganisms have wide applications in the agricultural industry as well as in the production of various secondary metabolites (Table 1) for human use, as described below.

### Agricultural applications

#### Plant growth promotion

It has been already reported that the mangrove microorganisms are beneficial for agriculture (Kathiresan and Selvam 2006). The rhizosphere soil of mangrove plants harbors a large number of beneficial bacteria with a large number of agricultural applications (Kathiresan and Selvam 2006). These strains have the ability to (1) fix nitrogen, (2) solubilize phosphate, (3) produce ammonia, and (4) produce the plant growth hormone indole acetic acid (IAA). Two strains identified from rhizosphere soil of mangroves viz., *Azotobacter vinelandii* and *Bacillus megaterium* have shown their ability to enhance mangrove seedlings (Kathiresan and Selvam 2006). Soil bacteria present in root regions are known to enhance plant growth. This beneficial effect is mediated through either direct or indirect mechanisms. The direct effects are commonly attributed to the supply of biologically fixed nitrogen and the production of plant hormones such as auxins. The indirect effects are suppression of bacterial, fungal and nematode pathogens, and production of siderophores, ammonia, antibiotics and volatile metabolites (Glick 1995). Further, it has been suggested that halophilic nitrifying bacteria can be used for removal of salinity and nitrate from waste water for recycling (Denariáz et al. 1989). The purified water could be used for irrigation of plants grown in arid soils. One of the problems with saline and hypersaline lands such as salt-affected soils is the relatively low microbial activity in these soils, which affects the vegetal and crop productivity of that soil. Therefore, the isolation of active bacteria from saline soil will allow the use of these bacteria in the reclamation of saline soils. The N<sub>2</sub>-fixing bacteria isolated from saline soil could be good candidates for use to improve the fertility of reclaimed arid and saline soils (Zahran et al. 1995). The inoculation of plants with plant-growth-promoting bacteria is a common tool in agriculture to enhance crop yields (Bashan and Holguin 1997).

#### Biopesticides

Natural marine products have the potential to replace chemical pesticides and other agents used to maximize crop yields and growth (Cardellina 1986). Nowadays, reports about mangrove fungi also reveal that many of them are able to produce insecticidal metabolites. According to Xiao et al. (2005), 188 marine-derived fungi were collected from

**Table 1** Natural products from mangrove microorganisms and their biotechnological applications

Natural product	Sources	Application	Reference
Xylanolytic enzyme	<i>Aspergillus niger</i> , <i>A. aculeatus</i> , <i>A. awamori</i> , <i>A. fumigates</i> , <i>Penicillium brasilianum</i> , <i>Chaetomium thermophile</i> , <i>Humicola insolens</i> , <i>Humicola</i> <i>lanuginosa</i> , <i>Humicola grisea</i> , <i>Melanocarpus albomyces</i> and <i>Bacillus</i> sp.	Paper manufacturing, animal feed, bread making, juice and wine industries and xylitol production	Polizeli et al. 2005
Phyllacoid, agar, carrageenan and alginate	<i>Codium fragile</i> , <i>C. latum</i> , <i>C. dwarkense</i> , <i>C. tomentosum</i>	Blood anticoagulant	Shanmugam and Mody 2000; Srivastava and Kulshreshtha 1989
Ethyl acetate extract, broth extract	Fungi	Biopesticide	Xiao et al. 2005; Chen et al. 1893
Polyketide synthases (PKSs)	Halophilic bacteria	Biosynthesis of secondary metabolites such as Erythromycin, Rapamycin, Tetracycline, Lovastatin and Resveratrol	Ventosa and Nieto 1995
Catalase, peroxidase, oxidase, polyphenol oxidase and ascorbic acid oxidase	<i>Pseudomonas aeruginosa</i> , <i>P.</i> <i>alcaligenes</i> , <i>Methylococcus</i> sp.	Industrial application	Mishra 2010
Amylase, protease, esterase and lipases	<i>Vibrionales</i> , <i>Actinomycetales</i> and <i>Bacillales</i>	Industrial application	Armando et al. 2009
Phytase	<i>Bacillus circulans</i> , <i>B. licheniformis</i> , <i>B. pantothenicus</i>	Industrial application	Joseph and Paul Raj 2007
Lipopeptides and glycolipids	<i>Bacillus subtilis</i> and <i>Pseudomonas</i> <i>aeruginosa</i>	Bio-surfactant	Pornsunthorntawee et al. 2008.
Aurantins A and B.	<i>Preussia aurantiaca</i>	Antimicrobial	Poch and Gloer 1989
Aigialomycins A-E, hypothemycin	Fungus, <i>Aigialus parvus</i> BCC 5311	Antimicrobial	Isaka et al. 2002
Secondary metabolites	Actinomycetes	Anti-tumor, anti-cancer and anti-infection properties	Hong et al. 2009
Xyloketal A (1), B (2), C (3), D (4), and E (5)	<i>Xylaria</i> sp.	Inhibitor of acetylcholine esterase (bioactive compound)	Lin et al. 2001
Isoflavone and prostaglandin analog	<i>Phomopsis</i> sp., <i>Paecilomyces</i> sp., <i>Sargassum</i> sp., <i>Halorosellinia</i> sp.	Anti-cancer	Tao et al. 2010
Poly aromatic hydrocarbon and oil degrading enzyme	<i>Pseudomonas</i> , <i>Shewanella</i> , <i>Sphingomonas</i> , <i>Arthrobacter</i> <i>Pseudomonas</i> , <i>Marinobacter</i> , <i>Alcanivorax</i> , <i>Microbulbifer</i> , <i>Sphingomonas</i> , <i>Micrococcus</i> , <i>Cellulomonas</i> , <i>Dietzia</i> and <i>Gordonia</i>	Bioremediation	Desai et al. 2010; Brito et al. 2006; Ramsay et al. 2000; Yu et al. 2005; Ke et al. 2003; Ventosa et al. 1998; Mishra et al. 2011
Laccase	<i>Basidiomycetes</i>	Bioremediation	D'souza et al. 2006; Sabu 2003
Phosphatase	<i>B. licheniformis</i> , <i>Chryseomonas</i> <i>luteola</i> , <i>Pseudomonas stutzeri</i> , <i>Aspergillus niger</i> etc.	Agriculture	Vazquez et al. 2000; Gyaneshwar et al. 1998
L-Asparaginase	<i>Halococcus</i>	Industrial	Sudha 1981
Xylanase, ligninase	<i>Aspergillus niger</i> and <i>Phlebia</i> sp.	Industrial	Li et al. 2002
Biosurfactant	<i>Leucobacter</i> , <i>B. subtilis</i> , <i>P. aeruginosa</i> ,	Industrial	Saimmai et al. 2011
Enniating G	<i>Fusarium</i> sp.	Antimicrobial	Lin et al. 2002a
Enalin A & B	<i>Verruculina enalia</i>	Antimicrobial	Lin et al. 2002b
2-Pyronone derivative	<i>Nocardia</i> sp.	Bioactive compound	Lin et al. 2001
Azapirones, P-terphenyles, shearinines D-K	<i>Penicillium</i> sp.	Bioactive compound	Huang et al. 2011; Xu et al 2007
$\beta$ -Carboline, adenosine, 8-hydroxyl-3, 5-dimethyl- isochroman-1-one	Fungi	Antitumor	Song et al. 2004

the sediment in Zhoushan Sea area, the mangrove at Yunxiao and Jiulongjiang estuary in Fujian Province, China, of which the ethyl acetate extract of strain 164 exhibited a strong lethal effect on nematode *Rhabditis* sp. Chen et al. (2003, 2006) isolated an endophytic fungus, *Kandelia candel* from an estuarine mangrove on the South China Sea coast, the broth extract of which exhibits cytotoxicity towards NCI4460 and Bel-7402, and high activities against *Heliothis armigera* (Hühner) and *Sinergasilus* sp. As the potentiality of mangrove-associated fungi in insecticidal activity has been investigated sporadically, continued research and development in this area will result in the discovery of new natural pesticides. The high potential of mangrove fungal metabolites has been anticipated as an alternative source of novel pesticide compounds. Also in Australia, *Bacillus thuringiensis*, which exhibits insecticidal activity against mosquito larvae of *Anopheles maculatus*, *Aedes aegypti* and *Culex quinquefasciatus*, has been isolated from mangrove sediments (Lee and Seleena 1990).

## Industrial applications

### Enzymes

Mangrove microorganisms have a diverse range of enzymatic activity and are capable of catalyzing various biochemical reactions with novel enzymes. Especially, halophilic microorganisms possess many hydrolytic enzymes (amylases, nucleases, phosphatases, and proteases) and are capable of functioning under conditions that lead to precipitation or denaturation of most proteins (Ventosa and Nieto 1995). It is believed that sea water, which is saline in nature and chemically closer to human blood plasma, could provide microbial products, in particular enzymes, that could be safer (i.e., having no or less toxicity or side effects) when used for therapeutic applications in humans (Sabu 2003). The hydrolase enzymes produced by halophilic bacteria are currently of commercial interest (Ventosa and Nieto 1995). Polyketide synthases (PKSs) are a class of enzymes involved in the biosynthesis of secondary metabolites such as Erythromycin, Rapamycin, Tetracycline, Lovastatin and Resveratrol, which have all now been isolated and purified from halophilic bacteria. A halophilic bacterium (*Halococcus*) producing L-asparaginase was also reported from a mangrove environment (Sudha 1981). From Brazil mangrove sediment, *Vibrionales* appeared to be the predominant enzyme-producing group within the community when compared with other groups (*Actinomycetales* and *Bacillales*), mainly for the production of amylase and protease (Dias et al. 2009). The order *Vibrionales* has been revealed to be metabolically versatile with a high production of enzymes. An isolate of *Vibrio fluvialis* from mangrove sediments was

used to produce an alkaline extracellular protease with high efficiency for use in industrial detergents (Venugopal and Saramma 2006). Mishra et al. (2010) evaluated and reported the activity of some stress enzymes such as catalase, peroxidase, oxidase, polyphenol oxidase and ascorbic acid oxidase from six Gram-negative bacteria isolated from mangroves of Bhitarkanika, Orissa, India. Bacteria isolated from mangrove sediment of Brazil were found to produce diverse extracellular enzymes such as amylase, protease, esterase and lipases (Dias et al. 2009). Joseph and Paul Raj (2007) reported five phytase-producing *Bacillus* strains from mangrove ecosystems of Kochin, Kerala, India. Three bacterial and one fungal strain producing tanase have been isolated from the mangrove forest of North Malabar, Kerala, India. Wu (1993) identified 15 genera (42 strains) of fungi from mangroves in the Tansui Estuary near Taipei, Taiwan, and found that most of the *ascomycetes* were able to secrete a wide range of enzymes potentially capable of decomposing mangrove litter. Raghukumar et al. (2004) reported that a mangrove fungus, *Aspergillus niger*, can produce a thermostable, cellulose-free alkaline xylanase that showed activity in biobleaching of paper pulp and its crude enzyme with high xylanase activity could bring about bleaching of sugarcane bagasse pulp by a 60-min treatment at 55°C. A marine hypersaline-tolerant white-rot fungus, *Phlebia* sp. MG-60, screened from mangrove stands (Li et al. 2002), has shown excellent lignin degrading ability. It can degrade more than 50% of lignin incubated with whole sugarcane bagasse and the whole sugarcane bagasse might be used to produce animal feed after fermentation (Li et al. 2003).

### Biosurfactants

Biosurfactants are getting much more attention compared to chemical surfactants owing to mild production conditions, lower toxicity, higher biodegradability and environmental compatibility (Mulligan 2009). All these qualities of biosurfactants have prompted their numerous applications in environmental protection as well as in the food, cosmetic, biopesticide and pharmaceutical industries (Singh and Cameotra 2004). Based on the types of biosurfactant-producing microbial species and the nature of their chemical structures, biosurfactants can be categorized into four main groups: lipopeptides or lipoproteins, glycolipids, phospholipids, and polymeric surfactants (Fathabad 2011). Among these four groups, the most common biosurfactants that have been isolated and studied are the lipopeptides produced by *Bacillus subtilis* strains, and the glycolipids produced by *Pseudomonas aeruginosa* strains (Pornsunthorntaweet et al. 2008). Biosurfactants have advantages over their chemical counterparts because they are bio-degradable, have low

toxicity, are effective at extreme temperatures or pH values and show better environmental compatibility (Mulligan 2009). Recently the *Leucobacter komagatae* 183 strain, isolated from mangrove sediment in Trang, southern Thailand, was evaluated as a potential biosurfactant producer (Saimmai et al. 2011).


## Pharmaceutical applications

### Antimicrobial compounds

The need for diversity and the development of new classes of antimicrobial compounds is increasing, due to trends in antibiotic resistance among different strains of bacteria, fungi and other microorganism, which are causing serious problems in the containment of infectious diseases. A significant number of reports have focused on antimicrobial metabolites isolated from mangrove saprophytic fungi. A fungal strain *Preussia aurantiaca* isolated from mangrove forest was found to synthesize two new despidones (Auranticins A and B) that display antimicrobial activity (Poch and Gloer 1991). Aigialomycins A–E, new 14-membered resorcylic macrolides, were isolated together with a known hypothemycin from the mangrove fungus, *Aigialus parvus* BCC 5311 (Isaka et al. 2002). Enniating G—a novel compound with a structure of cyclohexapeptide was also isolated from the culture broth of the mangrove fungus *Fusarium* sp., (Lin et al. 2002a; Lin and Zhou 2003; You et al. 2006), and displays antitumor, antibiotic, insecticidal and phytotoxic activity. An ascomycete, *Verruculina enalia*, is a common tropical species found on mangrove wood worldwide, reported to produce two new phenolic compounds, enalin A and B, with hydroxymethyl furfural and three cycloideptides from its fermentation broth. Enalin A is a coumaranone—a type of compound distributed widely from microorganisms to higher plants and having antimicrobial, antifungal, phytotoxic and antidiabetic activities (Lin et al. 2002b) Among the mangrove fungi, more and more mangrove endophytes now have been researched, and more and more antimicrobial metabolites have been isolated. One of them is Cytosporone B, which shows broad activities against fungi. In addition to fungi, Wiwin Retnowati (2010) reported an actinomycete *Streptomyces* sp., from mangrove soil in the eastern coast of Surabaya, Indonesia, capable of producing a series of antibiotics that strongly inhibit the growth of Gram-positive and Gram-negative bacteria. Santhi and Jebakumar (2011) reported some *Streptomyces* sp., from mangrove sediment of Manakudi estuary, India, exhibits potent antimicrobial effects against methicillin-resistant *Staphylococcus aureus* (clinical isolate) and methicillin-susceptible *S. aureus* and *Salmonella typhi*.

### Bioactive compounds


It is encouraging that bioactive compounds have been obtained from Mangrove plants, fungi, bacteria including actinomycetes (Cheng et al. 2009). Actinomycetes isolated from mangrove habitats are a potentially rich source for the discovery of anti-infection and anti-tumor compounds, and of agents for treating neurodegenerative diseases and diabetes (Hong et al. 2009). Mangrove ecosystems have been considered a “hot-spot” for newer and better drugs naturally produced by the microorganisms living in this environment. In the mangroves situated around the coast of China and surrounding islands, Hong et al. (2009) compiled more than 2,000 fungi-like bacteria or actinomycetes with the potential to synthesize biologically active secondary metabolites that conferred anti-tumor, anti-cancer and anti-infection properties. Using morphological, biochemical and molecular identification techniques and screening tests, approximately 20% showed activity against the growth of human colon tumor 116 cells, whereas only 3% inhibited the protein PTP1B associated with diabetes. Interestingly, it was discovered that, in general, most of the bioactive strains were found in plant tissues although more isolates were identified in rhizosphere soils. Lin et al. (2001) also reported three new 2-pyranon derivatives from the mangrove actinomycetes *Nocardioopsis* sp., A00203. Sivakumar et al. (2005) reported a *Streptomyces albidoflavus* from the pichavaram mangrove that showed antitumor properties. Five unique metabolites,



**Fig. 2** Five unique metabolites, xyloketals A (1), B (2), C (3), D (4), and E (5), and the unknown 6 were isolated from mangrove fungus *Xylaria* sp. (no. 2508), obtained from the South China Sea by Lin et al. (2001)

xyloketal A (1), B (2), C (3), D (4), and E (5) (Fig. 2) were isolated from mangrove fungus *Xylaria* sp. (no. 2508), obtained from the South China Sea. The structures of these compounds were elucidated by spectroscopic and X-ray diffraction experiments. Xyloketal A is a ketal compound with C3 symmetry and xyloketal B–E are its analogues. It was found that xyloketal C slowly rearranged to xyloketal B in DMSO-*d*<sub>6</sub> solution at room temperature. Xyloketal A exhibited the activity of inhibiting acetylcholine esterase (Lin et al. 2001). Similarly, Tao et al. (2010) reported Isoflavone and Prostaglandin analog compounds (Fig. 3) from mangrove fungi isolated from the South China Sea that appeared to be promising for treating cancer patients with multidrug resistance, which should encourage more efforts to isolate promising candidates for further development as clinically useful chemotherapeutic drugs from mangrove microorganisms. Huang et al. (2011) reported eight secondary metabolites (Fig. 4), including three new azaphilones (chermesinones A–C, 1–3), three new *p*-terphenyls (6'-*O*-desmethylterphenyllin, 4; 3-hydroxy-6'-*O*-desmethylterphenyllin, 5; 3''-deoxy-6'-*O*-desmethylcandidusin B, 7), and two known *p*-terphenyls (6, 8), were isolated from the culture of the mangrove endophytic fungus *Penicillium chermesinum* (ZH4-E2). Terphenyls 4, 5, and 6 exhibited strong inhibitory effects against  $\alpha$ -glucosidase and Terphenyls 7 and 8 showed inhibitory activity towards acetylcholinesterase.

Eight new indole triterpenes named shearinines D–K, along with shearinine A, paspalitrem A, and paspaline, have been isolated from the mangrove endophytic fungus *Penicillium* sp., Shearinines D, E, and (with reduced potency) G exhibit significant in vitro blocking activity on large-conductance calcium-activated potassium channels (Xu et al. 2007).  $\beta$ -Carboline, adenosine and 8-hydroxyl-3,5-dimethyl-isochroman-1-one, were isolated from mangrove fungus K32. The interaction of  $\beta$ -carboline with calf thymus DNA was investigated by UV-vis and fluorescence spectra, resulting in the occurrence of a binding reaction, which was proposed to be one possible mechanism of the antitumor activity of  $\beta$ -carboline (Song et al. 2004).




**Fig. 3** a Prostaglandin- and b isoflavone analogs isolated from mangrove fungi of South China Sea by Tao et al. (2010)

## Environmental applications

### Bioremediation

Bioremediation strategies can be improved by a greater knowledge of microbiology, diversity and function, and new molecular technologies can support this development. Genomics, for instance, provided complete genome sequence data for several microorganisms that are significant for bioremediation, such as *Pseudomonas*, *Shewanella*, *Sphingomonas*, *Arthrobacter* etc. (Desai et al. 2010). In addition to processing nutrients, mangrove bacteria may also help in processing industrial wastes. Iron-reducing bacteria are common in mangrove habitats in some mining areas (Panchnadikar 1993). Eighteen bacterial isolates that metabolize waste drilling fluid were collected from a mangrove swamp in Nigeria (Benka-coker and Olumagin 1995). The presence and activity of the oil-degrading microorganisms in mangrove sediments not only plays a key role in the bioremediation of oil in mangroves but is also considered to represent new prospects for the use of molecular tools to monitor bioremediation processes. The majority of oil from oceanic oil spills (e.g., the recent accident in the Gulf of Mexico) converges on coastal ecosystems such as mangroves. Microorganisms are directly involved in biogeochemical cycles as key drivers of the degradation of many carbon sources, including petroleum hydrocarbons. Various



**Fig. 4** Eight secondary metabolites, (1–3) azaphilones, (4–8) *p*-terphenyls [(4) 6'-*O*-desmethylterphenyllin, (5) 3-hydroxy-6'-*O*-desmethylterphenyllin, (7) 3''-deoxy-6'-*O*-desmethylcandidusin B] from the culture of the mangrove endophytic fungus *Penicillium chermesinum*, isolated from the South China Sea (Huang et al. 2011)

bacterial groups present in mangrove sediment, such as *Pseudomonas*, *Marinobacter*, *Alcanivorax*, *Microbulbifer*, *Sphingomonas*, *Micrococcus*, *Cellulomonas*, *Dietzia* and *Gordonia*, are already known for their capacity to degrade hydrocarbons (Brito et al. 2006). Bacteria were isolated in the sediment of a mangrove located in Hong Kong, China, which demonstrated a great capacity for polyaromatic hydrocarbon (PAH) degradation in vitro and could be used to degrade PAH in contaminated sediment (Ramsay et al. 2000). Similarly, Yu et al. (2005) investigated the biodegradability of PAHs, fluorine (Fl), phenanthrene (Phe) and pyrene by a bacterial consortium enriched with mangrove sediment. The consortium was formed by three bacterial strains: *Rhodococcus* sp., *Acinetobacter* sp., and *Pseudomonas* sp. Ramsay et al. (2000) reported a large number, and wide diversity, of PAH-degrading microorganisms in Australian mangrove sediments. In a study of contaminated mangrove microcosms, Ke et al. (2003) demonstrated the removal of 90 % of pyrene in 6 months. Some bacterial species, such as *Streptococcus*, *Staphylococcus*, *Micrococcus*, *Moraxella* and *Pseudomonas* and fungal species such as *Aspergillus glaucus* and *A. niger* were also reported that degrade polythene and plastic bags from the mangrove environment (Kathiresan 2003). D'Souza et al. (2006) reported a mangrove white-rot basidiomycetous fungus able to produce laccase to decolorize colored effluents and synthetic dyes. The efficiency of this fungus in decolorization of various effluents with laccase that is active at pH 3.0–6.0 and 60 °C in the presence of seawater has great potential in bioremediation of industrial effluent. Enhanced laccase production in the presence of industrial effluents in this fungus is an added advantage during bioremediation of effluents. Recently, two moderately halotolerant *Bacillus megaterium* species isolated from mangroves of Bhitarkanika, Orissa showed potential for reduction of toxic selenite to non-toxic elemental selenium (Mishra et al. 2011).

### Molecular methods for microbial taxonomy

Traditionally microbial taxonomy has been studied using a variety of morphological physiological and biochemical tests. However, the phenotypic characterization of bacteria is not sufficient to identify isolates beyond species level (Bakonyi et al. 2003). There also are problems associated with the study of bacterial and fungal diversity in soil due to methodological limitations and lack of taxonomic knowledge. Furthermore, the immense phenotypic and genetic diversity of soil bacteria and fungi make such studies very difficult. Less than 1 % of the bacterial populations found in nature are culturable, while 99 % are unculturable (Stanley 2002). Thus, many fungal populations are non-culturable.

Because of this limitation, bacterial biodiversity can be accurately determined only using molecular taxonomic tools that obviate the need for laboratory cultivation of isolates. The cultivation-independent method is called metagenomics or environmental genomics (also called ecogenomics), which is defined as the genomic analysis of microorganisms by direct extraction of nucleic acids from environmental samples (soil, water, sediment, etc.). It involves the amplification by PCR of DNA and cDNA from RNA extracted from environmental samples, and subsequent analysis of the diversity of the amplified molecules. Alternatively, the amplified products may be cloned and sequenced to identify and enumerate bacterial species present in the sample (Tasi and Olson 1991). A number of approaches have been developed to study molecular microbial diversity. These include DNA reassociation, DNA–DNA and mRNA:DNA hybridization, DNA cloning and sequencing, and other PCR-based methods, such as denaturing gradient gel electrophoresis (DGGE), temperature gradient gel electrophoresis (TGGE), ribosomal intergenic spacer analysis (RISA) and automated ribosomal intergenic spacer analysis (ARISA) (Muyzer 1999). FAME (fatty acid methyl ester) analysis, a biochemical method that does not rely on culturing of microorganisms, is also one of the methods to study microbial diversity.

### Future perspectives and concluding remarks

Microbial diversity in mangrove ecosystems is very rich and diverse, and has tremendous biotechnological potential. The study of the microbial diversity in mangrove is of vital importance to the understanding of the different processes of the mangrove, which may uncover potent novel microorganisms for screening of bioactive compounds. As evidenced from past and ongoing research, the microbial consortium has a plethora of bioactivity. A number of past reviews have focused the attention of researchers on this tremendous treasure of mangrove microbial diversity but there is still a long way to go. Interdisciplinary research and collaborative endeavors are required amongst scientists, microbiologists and biotechnologists to provide innovative approaches to study the microbial diversity and explore the biotechnological potential of this unique ecosystem. Using culture-dependent technologies reveals only a small percentage (<1 %) of the microbial community, leaving 99 % of microorganisms still undiscovered. Recent developments in molecular techniques will help to explore a good percentage of the microbial community that can be exploited for the welfare of the mankind.

## References

- Abdel-Wahab MAA (2005) Diversity of marine fungi from Egyptian Red Sea mangroves. *Bot Mar* 48:248–355
- Alias SA, Kuthubutheen AJ, Jones EBG (1995) Frequency of occurrence of fungi on wood in Malaysian mangroves. *Hydrobiologia* 295:97–106
- Alongi DM (1988) Bacterial productivity and microbial biomass in tropical mangrove sediments. *Microb Ecol* 15:59–79
- Alongi DM, Boto KG, Robertson AI (1992) Nitrogen and Phosphorus cycles in tropical mangrove ecosystems. Washington DC: Am Geophys Univ 41:251–292
- Alongi DM, Christoffersen P, Tirendi F (1993) The influence of forest type on microbial-nutrient relationships in tropical mangrove sediments. *J Exp Mar Biol Ecol* 171:201–223
- Alongi DM, Sasekumar A, Tirendi F, Dixon P (1998) The influence of stand age on benthic decomposition and recycling of organic matter in managed mangrove forests of Malaysia. *J Exp Mar Biol Ecol* 225:197–218
- Amann RI, Ludwig W, Schleifer KH (1995) Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiol Rev* 59:143–169
- Ara I, Kudo T, Matsumoto A, Takahashi Y, Omura S (2007) *Nonomuraea maheshkhaliensis* sp. nov., a novel actinomycete isolated from mangrove rhizosphere mud. *J Gen Appl Microbiol* 53:159–166
- Armando CFD, Andreote FD, Dini-Andreote F, Lacava PT, Sá ALB, Melo IS et al (2009) Diversity and biotechnological potential of culturable bacteria from Brazilian mangrove sediment. *World J Microbiol Biotechnol* 25:1305–1311
- Bakonyi T, Derakhshifar I, Grabensteiner L, Nowotny N (2003) Development and evaluation of PCR assays for the detection of *Paenibacillus* larvae in honey samples. Comparisons with isolation and biochemical characterization. *Appl Environ Microbiol* 69:1504–1510
- Bano N, Nisa MU, Khan N, Saleem M, Harrison PJ, Ahmed SI, Azam F (1997) Significance of bacteria in the flux of organic matter in the tidal creeks of the mangrove ecosystem of the Indus River Delta, Pakistan. *Mar Ecol Prog Ser* 157:1–12
- Bashan Y, Holguin G (1997) *Azospirillum* plant relationships: environmental and physiological advances. *Can J Microbiol* 43:103–121
- Benka-coker MO, Olumagin A (1995) Waste drilling fluid utilising microorganisms in a tropical mangrove swamp oilfield location. *Bioresour Technol* 53(3):211–215
- Borse BD (1988) Frequency of occurrence of marine fungi from Maharashtra coast, India. *Indian J Mar Sci* 17(2):165–167
- Brito EM, Guyoneaud R, Goñi-Urriza M, Ranchou-Peyruse A, Verbaere A, Crapez MAC (2006) Characterization of hydrocarbonoclastic bacterial communities from mangrove sediments in Guanabara Bay, Brazil. *Res Microbiol* 157:752–762
- Cardellina JH (1986) Marine natural products as leads to new pharmaceutical and agrochemical agents. *Pure Appl Chem* 58:365–374
- Chen GY, Lin YC, Wen L, Vrijmoed LLP, Jones EBG (1893) Two new metabolites of a marine endophytic fungus (No. 1893) from an estuarine mangrove on the South China Sea coast. *Tetrahedron* 59:4907–4909
- Chen GY, Lin YC, Vrijmoed LLP, Fong WF (2006) A new isochroman from the marine endophytic fungus 1893#. *Chem Nat Comp* 42:138–141
- Cheng ZS, Pan JH, Tang WC, Chen QJ, Lin YC (2009) Biodiversity and biotechnological potential of mangrove-associated fungi. *J For Res* 20(1):63–72
- Cordeiro-Marino M, Braga MRA, Eston VR, Fujii MT, Yokoya NA (1992) Mangrove macro algal communities in Latin America. The state of the art and perspectives. In: Seeliger U (ed) Coastal plant communities of Latin America. Academic, San Diego, pp 51–64
- Cribb AB, Cribb JW (1995) Marine fungi from Queensland-1. Papers Univ Queensland. Dept Bot 3:78–107
- Cundell AM, Brown MS, Stanford R, Mitchell R (1979) Microbial degradation of *Rhizophora mangle* leaves immersed in the sea. *Estuar Coast Mar Sci* 9:281–286
- D’Croz L, Del Rosario J, Holness R (1989) Degradation of red mangrove (*Rhizophora mangle* L.) leaves in the Bay of Panama. *Rev Biol Trop* 37:101–104
- D’Souza DT, Tiwari R, Sah AK, Raghukumar C (2006) Enhanced production of laccase by a marine fungus during treatment of colored effluents and synthetic dyes. *Enzym Microb Technol* 38:504–511
- Dar SA, Kleerebezem R, Stams AJM, Kuenen JG, Muyzer G (2008) Competition and coexistence of sulfate-reducing bacteria, acetogens and methanogens in a lab-scale anaerobic bioreactor as affected by changing substrate to sulfate ratio. *Appl Environ Microbiol* 78:1045–1055
- Das S, Lyla PS, Khan SA (2006) Spatial variation of aerobic culturable heterotrophic bacterial population in sediment of the Continental slope of western Bay of Bengal. *Indian J Mar Sci* 36(1):51–58
- Denariaz G, Payne WJ, Gall JLE (1989) A halophilic denitrifier *Bacillus halodnitrificans* sp. nov. *Int J Syst Bacteriol* 39:145–151
- Desai C, Pathak H, Madamwar D (2010) Advances in molecular and “omics” technologies to gauge microbial communities and bioremediation at xenobiotic/anthropogen contaminated sites. *Bioresour Technol* 101(6):1558–1569
- Dias ACF, Andreote FD, Dini-Andreote F, Lacava PT, Sa ALB, Melo IS, Azevedo JL, Araujo WL (2009) Diversity and biotechnological potential of culturable bacteria from Brazilian mangrove sediment. *World J Microbiol Biotechnol* 25(7):1305–1311
- Dias ACF, Andreote FD, Rigonato J, Fiore MF, Melo IS, Araujo WL (2010) The bacterial diversity in Brazilian non disturbed mangrove sediment. *Antonie Van Leeuwenhoek* 98:541–555
- Eccleston GP, Brooks PR, Kurtböke DI (2008) The occurrence of bioactive micromonosporae in aquatic habitats of the Sunshine Coast in Australia. *Mar Drugs* 6:243–261
- Fathabad EG (2011) Biosurfactants in pharmaceutical industry. *Am J Drug Discov Dev* 1:58–69
- Gadek PA (ed) (1998) Patch deaths in tropical Queensland rainforests: association and impact of *Phytophthora cinnamoni* and other soil borne pathogens. Cooperative Research Centre for Tropical Rain forest Ecology and Management, Technical Report, Cairns, 99 pp
- Garrettson-Cornell L, Simpson J (1984) Three new marine *Phytophthora* species from New South Wales. *Mycotaxon* 19:453–470
- Gayathri S, Saravanan D, Radhakrishnan M, Balagurunathan R, Kathiresan K (2010) Bioprospecting potential of fast growing endophytic bacteria from leaves of mangrove and salt-marsh plant species. *Indian J Biotechnol* 9:397–402
- Glick BR (1995) The enhancement of plant growth by free living bacteria. *Can J Microbiol* 41:109–117
- Gray JP, Herwig RP (1996) Phylogenetic analysis of the bacterial communities in marine sediments. *Appl Environ Microbiol* 62:4049–4059
- Gulve RM, Deshmukh AM (2011) Enzymatic activity of actinomycetes isolated from marine sediments. *Recent Res Sci Technol* 3(5):80–83
- Gupta N, Das S, Basak UC (2007) Use of extracellular activity of bacteria isolated from Bhitarkanika mangrove ecosystem of Orissa coast. *Malaysian J Microbiol* 3(2):15–18
- Gupta N, Mishra S, Basak UC (2009a) Microbial population in phyllosphere of mangroves grow in different salinity zones of Bhitarkanika (India). *Acta Bot Malacina* 34:1–5
- Gupta N, Mishra S, Basak UC (2009b) Diversity of *Streptomyces* in mangrove ecosystem of Bhitarkanika. *Iranian J Microbiol* 1(3):37–42
- Gupta N, Sahoo D, Basak UC (2010) Evaluation of in vitro solubilization potential of phosphate solubilising *Streptomyces* isolated from phyllosphere of *Heritiera fomes* (mangrove). *Afr J Microbiol Res* 4(3):136–142

- Gyaneshwar P, Naresh Kumar G, Parekh LJ (1998) Effect of buffering on the P-solubilizing ability of microorganisms. *World J Microbiol Biotechnol* 14:669–673
- Haight M (2005) Assessing the environmental burdens of anaerobic digestion in comparison to alternative options for managing the biodegradable fraction of municipal solid wastes. *Water Sci Technol* 52:553–559
- Hesse PR (1962) Phosphorus fixation in mangrove swamp muds. *Nature* 193:295–296
- Hicks BJ, Silvester WB (1985) Nitrogen fixation associated with the New Zealand mangrove *Avicennia marina* (Forsk) *Vierh.* Var. *resinifera* (Forst. F) Bakh. *Appl Environ Microbiol* 49: 955–959
- Holguin G, Guzman MA, Bashan Y (1992) Two new nitrogen fixing bacteria from the rhizosphere of mangrove trees: their isolation, identification and in vitro interaction with rhizosphere *staphylococcus* sp. *FEMS Microbiol* 101:207–216
- Holguin G, Bashan Y, Mendoza-Salgado RA, Amador E, Toledo G, Vazquez P, Amador A (1999) La Microbiología de los manglares. Bosques en la frontera entre el mar y la tierra. *Ciencia Desarrollo* 144:26–35
- Holguin G, Vazquez P, Bashan Y (2001) The role of sediment microorganisms in the productivity, conservation, and rehabilitation of mangrove ecosystems: an overview. *Biol Fertil Soil* 33:265–278
- Holmer M, Storkholm P (2001) Sulphate reduction and sulphur cycling in lake sediments: a review. *Freshw Biol* 46:431–451
- Hong K, Gao AH, Xie QY, Gao H, Zhuang L, Lin HP, Yu HP, Li J, Yao XC, Goodfellow M, Ruan JS (2009) Actinomycetes for marine drug discovery isolated from mangrove soils and plants in China. *Mar Drugs* 7:24–44
- Huang H, Lv J, Hu Y, Fang Z, Zhang K, Bao S (2008) *Micromonospora rifamycinica* sp. nov, a novel actinomycete from mangrove sediment. *Int J Syst Evol Microbiol* 58:17–20
- Huang H, Feng X, Xiao Z, Liu L, Li H, Ma L, Lu Y, Ju J, She Z, Lin Y (2011) Azaphilones and *p*-terphenyls from the mangrove endophytic fungus *Penicillium chermesinum* (ZH4-E2) isolated from the South China Sea. *J Nat Prod* 74(5):997–1002
- Hyde KD (1990) A comparison of the intertribal mycota of five mangrove tree species. *Asian Mar Biol* 7:93–108
- Hyde KD (1996) Marine fungi. In: Grunovic C, Mallett K (eds) *Fungi of Australia*, vol 1B. ABR/CSIRO, Canberra, pp 39–64
- Hyde KD, Jones EBG, Leano E, Pointing SB, Poonyth AD, Vrijmoed LLP (1998) Role of fungi in marine ecosystems. *Biodiv Conserv* 7:1147–1161
- Isaka M, Suyarnestakorn C, Tanticharoen M (2002) Aigialomycins A–E, new resorcylic macrolides from the marine mangrove fungus *Aigialus parvus*. *J Org Chem* 67:1561–1566
- Jones EBG, Abdel-Wahab MA (2005) Marine fungi from the Bahamas Islands. *Bot Mar* 48:356–364
- Joseph I, Paul Raj R (2007) Isolation and characterization of phytase producing *Bacillus* strains from mangrove ecosystem. *J Mar Biol Assoc India* 2:177–182
- Kathiresan K (2003) How do mangrove forests induce sedimentation. *Rev Biol Trop* 51:355–360
- Kathiresan K, Bingham BL (2001) Biology of mangroves and mangrove ecosystems. *Adv Mar Biol* 40:81–251
- Kathiresan K, Qasim SZ (2005) Biodiversity of mangrove ecosystems. Hindustan, New Delhi, p 51
- Kathiresan K, Selvam MM (2006) Evaluation of beneficial bacteria from mangrove soil. *Bot Mar* 49(1):86–88
- Ke L, Wang WQ, Wong TWY, Wong YS, Tam NFY (2003) Removal of pyrene from contaminated sediments by mangrove microcosms. *Chemosphere* 51:25–34
- Kohlmeyer J, Kohlmeyer E (1979) *Marine mycology. The higher fungi.* Academic, New York
- Kothamasi D, Kothamasi S, Bhattacharyya A, Kuhad RC, Babu CR (2006) Arbuscular mycorrhizae and phosphate solubilising bacteria of the rhizosphere of the mangrove ecosystem of Great Nicobar Island, India. *Biol Fertil Soils* 42:358–361
- Kristensen E, Holmer M, Bussarawit N (1991) Benthic metabolism and sulfate reduction in a south-east Asian mangrove swamp. *Mar Ecol Prog Ser* 73:93–103
- Lageiro MM, Moura MJ, Reis A, Ferreira MJC (2007) Microbial proteases application in leather industry. *J Biotechnol* 131:239–240
- Lakshmanaperumalsamy P (1987) Nitrogen fixing bacteria, *Azotobacter* sp. in aquatic sediment. *Fish Technol Soc Fish Technol* 24 (2):126–128
- Lakshmanaperumalsamy P, Chandramohan D, Natarajan R (1978) Antibacterial and antifungal activity of streptomycetes from Porto Novo coastal environment. *Mar Biol* 11:15–24
- Lee HL, Seleena P (1990) Effect of sodium chloride on the growth of several isolates of *Bacillus thuringiensis* Serotype H-14, isolated from mangrove swamp soil in Malaysia. *Mosquito Borne Dis Bull* 7(4):134–135
- Li X, Kondo R, Sakai K (2002) Biodegradation of sugarcane bagasse with marine fungus *Phlebia* sp. MG-60. *J Wood Sci* 48:159–162
- Li X, Kondo R, Sakai K (2003) Studies on hypersaline-tolerant white-rot fungi IV: effects of Mn and NH<sub>4</sub> on manganese peroxidase production and Roly R-478 decolorization by the marine isolate *Phlebia* sp. MG-60 under saline conditions. *J Wood Sci* 49:355–360
- Liang JB, Chen YQ, Lan CY, Tam FY, Zan QJ, Huang LN (2006) Recovery of novel bacterial diversity from mangrove sediment. *Mar Biol* 150:739–747
- Lin YC, Zhou SN (2003) Marine microorganism and its metabolites. Chemical Industry, Beijing, pp 426–427
- Lin YC, Wu XY, Feng S, Jiang GC, Luo JH, Zhou SN, Vrijmoed LLP, Jones EBG, Krohn K, Steingröver K, Zsila F (2001) Five unique compounds: xyloketals from mangrove fungus *Xylaria* sp. from the South China Sea coast. *J Org Chem* 66:6252–6256
- Lin YC, Wang J, Wu XY, Zhou SN, Vrijmoed LLP, Jones EBG (2002a) A novel compound enniatin G from the mangrove fungus *Halosarphaea* sp. (strain 732) from the South China Sea. *Aust J Chem* 55:225–227
- Lin YC, Wu XY, Deng ZJ, Wang J, Zhou SN, Vrijmoed LLP, Jones EBG (2002b) The metabolites of the mangrove fungus *Verrucolium enalia* No. 2606 from a salt lake in the Bahamas. *Phytochemistry* 59:469–471
- Liu A, Wu X, Xu T (2007) Research advances in endophytic fungi of mangrove. *Chin J Appl Ecol* 18(4):912–918
- Loka Bharathi PA, Oak S, Chandramohan D (1991) Sulfate-reducing bacteria from mangrove swamps II: their ecology and physiology. *Oceanol Acta* 14:163–171
- Lyimo TJ, Pol A, Op den Camp HJ (2002) Sulfate reduction and methanogenesis in sediments of Mtoni mangrove forest, Tanzania. *Ambio* 31:614–616
- Lyimo TJ, Pol A, Jetten SMM, Op den Camp HJM (2008) Diversity of methanogenic archaea in a mangrove sediment and isolation of a new *Methanococoides* strain. *FEMS Microbiol Lett* 291:247–253
- Mann FD, Steinke TD (1992) Biological nitrogen fixation (acetylene reduction) associated with decomposing *Avicennia marina* leaves in the Beach wood Mangrove Nature Reserve. *S Afr J Bot* 58:533–536
- Margesin R, Schinner F (2001) Potential of halotolerant and halophilic microorganisms for biotechnology. *Extremophiles* 5:73–83
- Maria GL, Sridhar KR (2002) Richness and diversity of filamentous fungi on woody litter of mangroves along the west coast of India. *Curr Sci* 83:1573–1580
- Maria GL, Sridha KR, Raviraja NS (2005) Antimicrobial and enzyme activity of mangrove fungi of south west coast of India. *J Agric Technol* 1:67–80
- Marty DG (1985) Description de quatre souches *Methanogenes* thermo tolerantis isolee de sediments marins ou intertidaux. *C R Acad Sci III* 300:545–548

- Matondkar SGP, Mahtani S, Mavinkurve S (1981) Studies on mangrove swamps of Goa, Heterotrophic bacterial flora from mangrove swamps. *Mahasagar Bull Natl Inst Oceanogr* 14:325–327
- Maxwell GS (1968) Pathogenicity and salinity tolerance of *Phytophthora* sp. isolated from *Avicennia resinifera* (Forst F.)—some initial investigations. *Tane* 14:13–23
- Mishra, RR (2010) Microbial biodiversity in mangroves of Bhitarkanika, Orissa—a study on genotypic, phenotypic and proteomic characterisation of some predominant bacteria. PhD thesis submitted to North Orissa University, Orissa, India
- Mishra RR, Prajapati S, Das J, Dangar TK, Das N, Thatoi HN (2011) Reduction of selenite to red elemental selenium by moderately halotolerant *Bacillus megaterium* strains isolated from Bhitarkanika mangrove soil and characterization of reduced product. *Chemosphere* 84(9):1231–1237
- Mitra A, Santra SC, Mukherjee J (2008) Distribution of actinomycetes and antagonistic behaviour with the physico-chemical characteristics of the world's largest tidal mangrove forest. *Appl Microbiol Biotechnol* 80:685–695
- Mobanraju R, Rajgopal BS, Daniels L, Natrajan R (1997) Isolation and characterisation of methanogenic bacteria from mangrove sediment. *J Mar Biotechnol* 5:147–152
- Mulligan CN (2009) Recent advances in the environmental applications of biosurfactants. *Curr Opin Colloid Interface Sci* 14:372–378
- Muyzer G (1999) DGGE/TGGE a method for identifying genes from natural ecosystems. *Curr Opin Microbiol* 2:317–322
- Muyzer G, Stams AJM (2008) The ecology and biotechnology of sulphate-reducing bacteria. *Nat Rev Microbiol* 6:441–454
- Nakagiri A, Newell SY, Ito T, Tan TK, Pek CL (1996) Biodiversity and ecology of the Oomycetous fungus, *Halophytophthora*. *DIWPA Series. Biodiversity and the Dynamics of Ecosystems* 1:273–280
- Nedumaran T, Thillairajsekhar K, Perumal P (2008) Mangrove associated cyanobacteria at Pichavaram, Tamilnadu. *Seaweeds Res Utiln* 30:77–85
- Nedwell DB, Blackburn TH, Wiebe WJ (1994) Dynamic nature of the turnover of organic carbon, nitrogen and sulphur in the sediments of a Jamaican mangrove forest. *Mar Ecol Prog Ser* 110:223–231
- Odum WE, Heald EJ (1972) Trophic analyses of an estuarine mangrove community. *Bull Mar Sci* 22:671–738
- Odum WE, Heald EJ (1975a) Mangrove forests and aquatic productivity. In: Hasler AD (ed) *Coupling of land and water systems. Ecological studies series*. Springer, Berlin, pp 129–136
- Odum WE, Heald EJ (1975b) The detritus-based food web of an estuarine mangrove community. In: Ronin LT (ed) *Estuarine research*. Academic, New York, pp 265–286
- Oremland RS, Marsh LM, Polcin S (1982) Methane production and simultaneous sulfate reduction in anoxic salt marsh sediments. *Nature* 296:143–145
- Pal AK, Purkayastha RP (1992) New Parasitic fungi from Indian mangrove. *J Mycopathol Res* 30(2):173–176
- Panchnadikar VV (1993) Studies of iron bacteria from mangrove ecosystem in Goa and Konkan. *Int J Environ Stud* 45(1):17–21
- Pegg KG, Gillespie NC, Forsberg LI (1980) *Phytophthora* spp. associated with mangrove death in central coastal Queensland. *Australas Plant Pathol* 9:6–7
- Pelegri SP, Twilley RR (1998) Heterotrophic nitrogen fixation (acetylene reduction) during leaf litter decomposition of two mangrove species from South Florida, USA. *Marine Biol* 131(1):53–61
- Poch GK, Gloer JB (1989) Helicascolides A and B: new lactones from the marine fungus *Helicascus kunaloanus*. *J Nat Prod* 52:257–260
- Poch GK, Gloer JB (1991) Auranticins A and B: two depsidones from a mangrove isolate of the fungus *Preussia aurantiaca*. *J Nat Prod* 54:213–217
- Polizeli MLTM, Rizzatti ACS, Monti R, Terenzi HF, Jorge JA, Amorim DS (2005) Xylanases from fungi: properties and Industrial applications. *Appl Microbiol Biotechnol* 67:577–591
- Pornsunthorntawe O, Wongpanit P, Chavadej S, Abe M, Rujiravanit R (2008) Structural and physicochemical characterization of crude biosurfactants produced by *Pseudomonas aeruginosa* SP4 isolated from petroleum-contaminated soil. *Bioresource Technol* 99:1589–1595
- Prabhakaran NR, Gupta N, Krishnankutty M (1990) Fungal activity in Mangalvan—an estuarine mangrove ecosystem. In: Nair NB (ed) *Proceedings of the National Seminar on Estuarine Management, Trivandrum*. Academic, New York, pp 458–463
- Raghukumar S, Sathe-Pathak V, Sharma S, Raghukumar C (1995) Thraustochytrid and fungal component of marine detritus. *Field studies on decomposition of leaves of the mangrove Rhizophora apiculata*. *Aquat Microb Ecol* 9:117–125
- Raghukumar C, Muraleedharan U, Gaud VR, Mishra R (2004) Xylanases of marine fungi of potential use of bioleaching of paper pulp. *J Ind Microbiol Biotechnol* 31:433–441
- Ramachandran S, Venugopalan VK (1987) Nitrogen fixation by blue green algae in porto Novo Marine environment. *J Mar Biol Ass India* 29(1–2):337–343
- Ramamurthy T, Raju RM, Natarajan R (1990) Distribution and ecology of methanogenic bacteria in mangrove sediments of Pichavaram, east coast of India. *Indian J Mar Sci* 19:269–273
- Ramanathan AL, Singh G, Majumdar J, Samal AC, Chauhan R, Ranjan RK, Rajkumar K, Santra SC (2008) A study of microbial diversity and its interaction with nutrients in the sediments of Sundarban mangroves. *Indian J Mar Sci* 37(2):159–165
- Ramsay MA, Swannell RPJ, Shipton WA, Duke NC, Hill RT (2000) Effect of bioremediation community in oiled mangrove sediments. *Mar Pollut Bull* 41:413–419
- Ravikumar S (1995) Nitrogen fixing *Azotobacters* from the mangrove habitat and their utility as biofertilizers. PhD thesis, Annamalai University, India
- Ravikumar DR, Vittal BPR (1996) Fungal diversity on decomposing biomass of mangrove plant *Rhizophora* in Pichagram estuary, east coast of India. *Indian J Mar Sci* 21(1):64–66
- Ravikumar S, Fredimoses M, Gokulakrishnan R (2011a) Biodiversity of actinomycetes in Manakkudi mangrove ecosystem, Southwest coast of India. *Ann Biol Res* 2(1):76–82
- Ravikumar S, Inbaneson SJ, Uthiraselvam M (2011b) Priya SR2, Ramu A, Banerjee MB (2010) Diversity of endophytic actinomycetes from Karangkadu mangrove ecosystem and its antibacterial potential against bacterial pathogens. *J Pharm Res* 4(1):294–296
- Rivera-Monroy VH, Day WJ, Twilley RR, Vera-Herrera F, Coronado-Molina C (1995a) Flux of nitrogen and sediment in a fringe mangrove forest in Terminos lagoon, Mexico. *Estuar Coast Shelf Sci* 40:139–160
- Rivera-Monroy VH, Twilley RR, Boustany RG, Day WJ, Vera-Herrera F, Ramirez MC (1995b) Direct denitrification in mangrove sediments in Términos Lagoon, Mexico. *Mar Ecol* 126:97–109
- Roy S, Hens D, Biswas D, Biswas D, Kumar R (2002) Survey of petroleum—degrading bacteria in coastal waters of Sunderban biosphere reserve. *World J Microbiol Biotechnol* 18:575–581
- Sabu A (2003) Sources, properties and applications of microbial therapeutic enzymes. *Ind J Biotechnol* 2(3):334–341
- Sadaba RB, Vrijmoed LLP, Jones EBG, Hodgkiss IJ (1995) Observations on vertical distribution of fungi associated with standing senescent *Acanthus ilicifolius* stems at Mai Po mangrove, Hong Kong. *Hydrobiologia* 295:119–126
- Saho K, Dhal NK (2009) Potential microbial diversity in mangrove ecosystem: a review. *Indian J Mar Sci* 38(2):249–256
- Sahu MK, Sivakumar K, Kannan L (2005) Isolation of actinomycetes from different samples of the Vellar estuary, southeast coast of India. *Pollut Res* 24:45–48
- Sahu MK, Sivakumar K, Thangaradjou T, Kannan L (2007) Phosphate solubilizing actinomycetes in the estuarine environment: an inventory. *J Environ Biol* 28:795–798

- Saimmai A, Sobhon V, Maneerat S (2011) Production of biosurfactants from a new and promising strain of *Leucobacter komagatae* 183. *Ann Microbiol* 62(1):391–402
- Santhi VS, Jebakumar SRD (2011) Phylogenetic analysis and antimicrobial activities of *Streptomyces* isolates from mangrove sediment. *J Basic Microbiol* 51:71–79
- Santos HF, Cury JC, Carmo FL, Santos AL, Tiedje J, Elsas JD, Rosado SA, Peyote RS (2011) Mangrove bacterial diversity and the impact of oil contamination revealed by pyrosequencing: bacterial proxies for oil pollution. *PLoS One* 6(3):e16943
- Sarma VV, Hyde KD (2001) A review on frequently occurring fungi in mangrove. *Fung Divers* 8:1–34
- Saxena D, Loka-Bharathi PA, Chandramohan D (1988) Sulfate reducing bacteria from mangrove swamps of Goa, central west coast of India. *Indian J Mar Sci* 17:153–157
- Sen N, Naskar K (2003) *Algal flora of Sundarbans Mangal*. Daya, New Delhi
- Sengupta A, Chaudhuri S (2002) Arbuscular mycorrhizal relations of mangrove plant community at the Ganges river estuary in India. *Mycorrhiza* 12:169–174
- Sengupta A, Choudhury S (1990) Halotolerant *Rhizobium* Strains from mangrove swamps of the Ganges River Delta. *Indian J Microbiol* 30(4):483–484
- Sengupta A, Choudhury S (1991) Ecology of Heterotrophic dinitrogen fixation in the rhizosphere of mangrove plant community at the Ganges river estuary in India. *Oecologia* 87:560–564
- Shanmugam S, Mody KH (2000) Heparinid active sulphated polysaccharides from marine algae as potential blood coagulant agents. *Curr Sci* 79:1672–1683
- Shearer CA, Descals E, Kohlmeyer B, Kohlmeyer J, Marvanova L, Padgett D, Porter D, Raja HA, Schmit JP, Thornton HA, Voglymayr H (2007) Fungal diversity in aquatic habitats. *Biodivers Conserv* 16:49–67
- Sherman RE, Fahey TJ, Howarth RW (1998) Soil-plant interactions in a neotropical mangrove forest: iron, phosphorus and sulfur dynamics. *Oecologia* 115:553–563
- Shoreit AAM, EL- Kady IA, Sayed WF (1994) Isolation and identification of purple non sulphur bacteria of mangal and non-mangal vegetation of red sea coast, Egypt. *Limnologia* 24:177–183
- Singh P, Cameotra SS (2004) Potential applications of microbial surfactants in biomedical sciences. *Trends Biotechnol* 22:142–146
- Sivakumar K (2001) Actinomycetes of an Indian mangrove (Pichavaram). Environment. PhD thesis, Annamalai University
- Sivakumar K, Sahu MK, Kathiresan K (2005) An antibiotic producing marine *Streptomyces* from the Pichavaram mangrove environment. *Journal of the Annamalai University, Part-B* XLI:9–18
- Sivakumar K, Sahu MK, Thangaradjou T, Kannan L (2007) Research on marine *Actinobacteria* in India. *Ind J Microbiol* 47:186–196
- Sivaramakrishnan S, Gangadharan D, Nampoothiri KM, Soccol CR, Pandey A (2006)  $\alpha$ -Amylases from microbial sources—an overview on recent developments. *Food Technol Biotechnol* 44:173–184
- Song XH, Liu XH, Lin YC (2004) Metabolites of mangrove fungus No. K23 and interaction of carboline with DNA. *J Trop Oceanogr* 23(3):66–71.
- Srivastava R, Kulshreshtha DK (1989) Bioactive polysaccharides from plants. *Phytochemistry* 28:2877–2883
- Stanley J (2002) *Biodiversity of Microbial life*. Wiley-Liss, New York
- Steinbüchel A, Fuchtenbusch B (1998) Bacteria and other biological systems for polyester production. *Trends Biotechnol* 16:419–427
- Steinke TD, Barnabas AD, Somaru R (1990) Structural changes and associated microbial activity accompanying decomposition of mangrove leaves in Mgeni Estuary. *S Afr J Bot* 56:39–48
- Sudha V (1981) Studies (a) on halophilic bacteria associated with mangrove sediment and a biovalve mollusc *Anadara rhombea* (Born) (Arcidae) and (b) on L-asparaginase (Anti-Leukemic agent) from an extremely halophilic bacterium. PhD Thesis. Annamalai University, Parangipettai, India
- Sundararaj V, Dhevendran K, Chandramohan D, Krishnamurthy K (1974) Bacteria and primary production. *Indian J Mar Sci* 3:139–141
- Suryanarayanan TS, Kumaresan V, Johnson JA (1998) Foliar fungal endophytes from two species of the mangrove *Rhizophora*. *Microbiology* 44:1003–1006
- Tabao NC, Moasalud RG (2010) Characterisation and identification of high cellulose-producing bacterial strains from Philippine mangroves. *Philipp J System Biol* 4:13–20
- Taketani GR, Yoshiura AC, Dias FCA, Andreote DF, Tsai MS (2010) Diversity and identification of methanogenic archaea and sulphate-reducing bacteria in sediments from a pristine tropical mangrove. *Antonie van Leeuwenhoek* 97:401–411
- Tao L, Zhang JY, Liang YJ, Chen LM, Zhen LC, Wang F, Mi Y, She ZG, To KKW, Lin YC, Fu LW (2010) Anticancer effect and structure-activity analysis of marine products isolated from metabolites of mangrove fungi in the South China Sea. *Mar Drugs* 8:1094–1105
- Tasi-Li-Yu, Olson BH (1991) Rapid method for direct extraction of DNA from soil and sediments. *Appl Environ Microbiol* 57:1070–1074
- Taylor LE, Henrissat B, Coutinho PM, Ekborg NA, Hutcheson SW, Weiner RM (2006) Complete cellulase system in the marine bacterium *Saccharophagus degradans* strain 2-40 T. *J Bacteriol* 188:3849–3861
- Thatoi HN, Biswal AK (2008) Mangroves of Orissa coast: floral diversity and conservation status. Special habitats and threatened plants of India. *ENVIS Wild Life And Protected Area* 11 (1):201–207
- Toledo G, Bashan Y, Soeldner A (1995) Cyanobacteria and black Mangroves in North Western Mexico. Colonization and diurnal and seasonal nitrogen fixation on aerial roots. *Can J Microbiol* 41:999–1011
- Urakawa H, Kita-Tsukamoto K, Ohwada K (1999) Microbial diversity in marine sediments from Sagami Bay and Tokyo Bay, Japan, as determined by 16S rRNA gene analysis. *Microbiology* 145:3305–3315
- Vassileva M, Azcon R, Barea Miguel J, Vassilev N (1998) Application of an encapsulated filamentous fungus in solubilisation of inorganic phosphate. *J Biotechnol* 63(1):67–72
- Vazquez P, Holguin G, Puerte ME, Lopez-Cortes A, Bashan Y (2000) Phosphate-solubilising micro organisms associated with the rhizosphere of mangroves in a semiarid Coastal lagoon. *Biol Fertil Soils* 30:460–468
- Venkateswara Sarma V, Hyde KD, Vittal BPR (2001) Frequency of occurrence of mangrove fungi from the east coast of India. *Hydrobiologia* 455:41–53
- Venkateswaran K, Natarajan R (1983) Seasonal distribution of inorganic phosphate solubilising bacteria and phosphatase producing bacteria in Porto Novo waters. *Indian J Mar Sci* 12(4):213–217
- Ventosa A, Nieto JJ (1995) Biotechnological applications and potentialities of halophilic microorganisms. *World J Microbiol Technol* 11:85–94
- Ventosa A, Nieto JJ, Oren A (1998) Biology of moderately halophilic bacteria. *Microbiol Mol Biol Rev* 62:504–544
- Venugopal M, Saramma AV (2006) Characterization of alkaline protease from *Vibrio fluvialis* strain VM10 isolated from a mangrove sediment sample and its application as a laundry detergent additive. *Process Biochem* 41:1239–1243
- Vethanayagam RR (1991) Purple photosynthetic bacteria from a tropical mangrove environment. *Mar Biol* 110:161–163
- Vethanayagam RR, Krishnamurthy K (1995) Studies on anoxygenic photosynthetic bacterium *Rhodospseudomonas* spp. from the tropical mangrove environment. *Ind J Mar Sci* 24:19–23
- Vrijmoed LLP, Jones EBG, Hyde KD (1991) Observations on subtropical mangrove fungi in the Pearl River Estuary. *Acta Sci Nat* 33(1):78–85
- Wang GW, Li HY, Sun WB (2003) Primary study on arbuscular mycorrhizas of mangrove in Qinzhou Bay. *Guihaia* 23(5):445–449

- Wiwini R (2010) Identification of *Streptomyces* sp-MWS1 producing antibacterial compounds. Indonesian J Trop Infect Dis 1(2):80–85
- Wu RY (1993) Studies on the microbial ecology of the Tansui Estuary. Bot Bull Acad Sin 34:13–30
- Xiao YT, Zheng ZH, Huang YJ, Xu QY, Su WJ, Song SY (2005) Nematicidal and brine shrimp lethality of secondary metabolites from marine-derived fungi. Journal of Xiamen University (Nature Science) 44(6):847–850
- Xie XC, Mei WL, Zhao YX, Hong K, Dai HF (2006) A new degraded sesquiterpene from marine actinomycete *Streptomyces* sp. 0616208. Chin Chem Lett 17:1463–1465
- Xu MJ, Gessner G, Groth I, Lange C, Christner A, Bruhn T, Deng ZW, Li X, Heinemann SH, Grabley S, Bringmann G, Sattler I, Lin WH (2007) Shearing D–K, new indole triterpenoids from an endophytic *Penicillium* sp. (strain HKI0459) with blocking activity on large-conductance calcium-activated potassium channels. Tetrahedron 63:435–444
- Yakimov MM, Abraham WR, Meyer H, Laura G, Golyshin PN (1999) Structural characterisation of lichenycin. A component by fast atom bombardment tandem mass spectrometry. Biochem Biophys Acta 1438:230–280
- You JL, Mao W, Zhou SN, Wang J, Lin YC, Wu SY (2006) Fermentation conditions and characterization of endophytic fungus #732 producing novel enniatin G from South China Sea. Act Sci Nat 45(4):75–78
- Yu KS, Wong AH, Yau KW, Wong YS, Tam NF (2005) Natural attenuation, bio stimulation and bioaugmentation on biodegradation of polycyclic aromatic hydrocarbons (PAHs) in mangrove sediments. Mar Pollut Bull 51:1071–1077
- Zahran HH, Ahmad MS, Afkar EA (1995) Isolation and characterization of nitrogen-fixing moderate halophilic bacteria from saline soils of Egypt. J Basic Microbiol 35:269–275
- Zhang YHP, Lynd LR (2004) Kinetics and relative importance of phosphorolytic and hydrolytic cleavage of cellodextrins and cellobiose in cell extracts of *Clostridium thermocellum*. Appl Environ Microbiol 70:1563–1569
- Zuberr DA, Silver WS (1978) Biological dinitrogen fixation (acetylene reduction) associated with florida mangroves. Appl Environ Microbiol 35:567–575