

UNIVERSITÀ DEGLI STUDI DI MILANO

# **ORIGINAL ARTICLE**



# Phylogenetically diverse bacteria isolated from tattoo inks, an azo dye-rich environment, decolorize a wide range of azo dyes



Seong Won Nho<sup>1+</sup>, Xuewen Cui<sup>1,2+</sup>, Ohgew Kweon<sup>1+</sup>, Jinshan Jin<sup>1+</sup>, Huizhong Chen<sup>1</sup>, Mi Sun Moon<sup>3</sup>, Seong-Jae Kim<sup>1\*</sup> and Carl E. Cerniglia<sup>1\*</sup>

# Abstract

**Purpose:** There has been an interest in the microbial azo dye degradation as an optional method for the treatment of azo dye-containing wastes. Tattoo ink is an extremely unique azo dye-rich environment, which have never been explored in terms of microorganisms capable of degrading azo dyes. Previously, we isolated 81 phylogenetically diverse bacteria, belonging to 18 genera and 52 species, contaminated in tattoo inks. In this study, we investigated if these bacteria, which can survive in the azo dye-rich environment, have an ability to degrade azo dyes.

**Methods:** We conducted a two-step azo dye degradation (or decolorization) assay. In step 1, a high-throughput degradability assay was done for 79 bacterial isolates using Methyl Red and Orange II. In step 2, a further degradation assay was done for 10 selected bacteria with a representative of 11 azo dyes, including 3 commercial tattoo ink azo dyes. Degradation of azo dyes were calculated from measuring optical absorbance of soluble dyes at specific wavelengths.

**Results:** The initial high-throughput azo dye assay (step 1) showed that 79 isolates had a complete or partial degradation of azo dyes; > 90% of Methyl Red and Orange II were degraded within 24 h, by 74 and 20 isolates, respectively. A further evaluation of azo dye degradability for 10 selected isolates in step 2 showed that the isolates, belonging to *Bacillus, Brevibacillus, Paenibacillus*, and *Pseudomonas*, exhibited an excellent decolorization ability for a wide range of azo dyes.

**Conclusions:** This study showed that phylogenetically diverse bacteria, isolated from azo dye-rich tattoo inks, is able to degrade a diverse range of azo dyes, including 3 azo dyes used in commercial tattoo inks. Some of the strains would be good candidates for future studies to provide a systematic understanding of azo dye degradation mechanisms.

Keywords: Tattoo ink, Microbial contamination, Bacterial azo dye degradation

equally to this work. <sup>1</sup>Division of Microbiology, National Cent

<sup>1</sup>Division of Microbiology, National Center for Toxicological Research, U.S. Food and Drug Administration, Jefferson, AR, USA

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

<sup>\*</sup> Correspondence: seong-jae.kim@fda.hhs.gov; carl.cerniglia@fda.hhs.gov \*Seong Won Nho, Xuewen Cui, Ohgew Kweon and Jinshan Jin contributed

## Background

Azo dyes, which are the most common class of dyes containing one or more azo (-N=N-) bonds, are widely used in textile, paper, cosmetic, pharmaceutical, and food industries (Carmen and Daniela 2012). Reports in the literature indicate that azo dyes and their intermediates may be toxic or mutagenic and carcinogenic, posing negative impact on the environmental health (Chung 2016; Feng et al. 2012). Industries release large amount of insufficiently treated azo dye-containing waste in the environment (Rawat et al. 2016).

Physicochemical-based methods, such as absorption, coagulation, precipitation, chemical transformation, and incineration, have often been used for the remediaion of azo dyes-containing waste effluents in the environment (Robinson et al. 2001). However, these methods have limited applicability due to lack of efficiency, high cost, generation of toxic by-products, and high energy requirements. In recent years, biotechnological methods using microorganisms have been suggested as an option to overcome the limitations of the physicochemical-based approaches for the reduction of azo dyes in the environment (Forgacs et al. 2004; Robinson et al. 2001). Diverse categories of microorganisms, including bacteria, fungi, and algae, have been reported to be capable of degrading a wide range of azo dyes (Solís et al. 2012; Stolz 2001).

Previously, we isolated upwards of 80 bacteria, belonging to 18 genera and 52 species, from a survey study of unopened sealed bottles of tattoo inks for microbial composition (Nho et al. 2020; Nho et al. 2018). Although the survey study was originally initiated to determine if microbial contaminants occurred in the tattoo inks, results of the study led us into a different question; that is, whether the presence of such microorgnaisms has something further to do with tattoo inks, an environment known to be highly abundant with azo dyes (Bäumler et al. 2000). We therefore decided to study whether there was a functional connection between the isolates found in the tattoo inks, which can survive in an azo dye-rich environment, and their ability to biodegrade azo dyes. Here, we present the data on the azo dye degradation (or decolorization) assay that we conducted on these phylogenetically diverse bacterial isolates.

Azo dye-degrading microorganisms have often been isolated from various environments, particularly including textile effluents, activated sludge, or azo dyecontaminated soils and ground waters near textile industries (Gomare and Govindwar 2009; Jin et al. 2007; Meehan et al. 2001; Srinivasan et al. 2014; Telke et al. 2010; Xu et al. 2005). However, to our knowledge, tattoo inks, an extremely unique azo dye-rich environment, have previously never been explored in terms of azo dye degradation. In this study, we examined if these bacterial isolates have an ability of azo dye degradation, and if so, how well can they degrade structurally different kinds of azo dyes.

## Materials and methods Dyes and chemicals

Methyl Red, Sudan I, Sudan II, Sudan IV, Toluidine Red, Orange II, Fast Dark Blue R Salt, Lithol Rubin BK, Orange G, Amaranth, Ponceau BS, Direct Blue 15, and dimethyl sulfoxide (DMSO) were purchased from Sigma-Aldrich Inc. (Saint Louis, MO, USA). Solvent Red 1, Hansa Yellow, Hansa Orange, and Alphamine Red R were purchased from Fisher Scientific (Hanover Park, IL, USA). Diarylide Yellow HR (Pigment Yellow 83) was purchased from Santa Cruz Biotechnology Inc. (Dallas, TX, USA). Chemical structures of azo dyes used in this study were shown in Fig. 1. Tryptic Soy Agar (TSA) and Tryptic Soy Broth (TSB) used for bacterial growth were purchased from Fisher Scientific (Hanover Park, IL, USA). Stock solutions of azo dyes (10 mM) were prepared by using 100% DMSO as solvent, then dye solutions were diluted with TSB in degradation assay. Final concentration of DMSO and dye in the assays was 1% and 100 µM except as indicated otherwise.

# Bacterial growth conditions and azo dye decolorization assay using 96-well microphates

All 79 bacterial strains used in this study (Table 1) were isolated from tattoo inks in our previous study (Nho et al. 2018). Bacteria were incubated overnight in TSB at 37 °C with shaking at 250 rpm. The overnight grown bacterial cultures were inoculated into 96-well microplates containing TSB with azo dyes or with 1% DMSO as carrier control. About  $5 \times 10^7 - 1 \times 10^8$  bacterial cells were inoculated into each well and microplates were incubated at 37 °C without shaking. Optical absorbance was measured for different dyes at specific wavelength (Table 1) at different time points (1 h, 2 h, 4 h, 6 h, 8 h, 18 h, and 24 h) after inoculation using Spectra ax M2 plate reader (Molecular Devices, USA). Calibration curve was determined by measuring optical absorbance of TSB with known concentrations of azo dyes in 96-well microplates. The amount of azo dyes remained in culture media was determined by measuring optical absorbance and calculation with calibration curve. The degradation (%) of azo dyes = 100 - (amount of remain azo dyes/ start amount of azo dye)  $\times$  100. This experiment was triplicated and repeated twice. The final decolorization (%) is the mean of data from repeated triplicated assay.

# Azo dye decolorization assay using 15-mL conical centrifuge tubes

Selected 10 bacterial species (Table 2) were grown overnight in TSB at 37  $^{\circ}$ C with shaking at 250 rpm. About 1



 $\times 10^8$  bacterial cells from overnight culture were inoculated into 15-mL conical centrifuge tubes containing 2-mL TSB with azo dye or with 1% DMSO as carrier control, then they were incubated at 37 °C for 48 h without shaking. Bacterial cultures were collected at 48 h by centrifugation at 10,000 rpm for 30 min with Dorval RC6+ Centrifuge (Thermo Scientific). Centrifugation was repeated one more time for the supernatant. Then the supernatants were subjected for measuring optical absorbance of soluble azo dyes at specific wave length (Table 1). To extract insoluble azo dyes, centrifugation pellets were re-suspended with 100% DMSO, and then

centrifuged at 10,000 rpm for 10 min. The supernatants of DMSO resuspension were subjected for measuring optical absorbance of insoluble azo dyes. Calibration curve was determined by measuring optical absorbance of known concentrations azo dyes in presence of TSB or 100% DMSO. The amount of soluble azo dyes was calculated with calibration curve determined in the presence of TSB. The amount of insoluble azo dyes was calculated with calibration curve determined in the presence of 100% DMSO. The degradation (%) of azo dyes = 100 – [(amount of soluble azo dye + amount of insoluble azo dye)/initial amount of azo dye]  $\times$  100. This

# Table 1 Degradation (%) of Methyl Red and Orange II by 79 bacterial isolates<sup>a</sup>

Isolate	Identity based on 16S rRNA	BLAST on NCBI 16S RefSeq database		Methyl	Red <sup>b</sup>		Orange	e II <sup>b</sup>	
#		Type strain (Sequence ID)	% Identity	1 h	8 h	24 h	1 h	8 h	24 h
1	Paenalcaligenes suwonensis	P. suwonensis ABC02-12 (NR_133804.1)	99.23	20 ± 3	96 ± 4	100 ± 0	5 ± 3	18 ± 5	92 ± 8
4	Bacillus acidiceler	B. acidiceler CBD 119 (NR_043774.1)	99.67	99 ± 1	$100 \pm 0$	$100 \pm 0$	9 ± 1	$25 \pm 4$	$43 \pm 3$
5	Bacillus velezensis	B. velezensis FZB42 (NR_075005.2)	99.64	74 ± 10	$100 \pm 0$	$100 \pm 0$	8 ± 2	12 ± 1	25 ± 2
6	Bacillus velezensis	B. velezensis FZB42 ( NR_075005.2)	99.74	$100 \pm 0$	$100 \pm 0$	$100 \pm 0$	2 ± 1	8 ± 4	24 ± 2
7	Bacillus haikouensis	B. haikouensis C-89 ( NR_148273.1)	99.39	62 ± 13	$100 \pm 0$	$100 \pm 0$	7 ± 2	13 ± 2	$62 \pm 5$
8	Bacillus marisflavi	B. marisflavi TF-11 (NR_025240.1)	99.73	$100 \pm 0$	$100 \pm 0$	$100 \pm 0$	$5 \pm 1$	$13 \pm 1$	$38 \pm 3$
9	Bacillus drentensis	B. drentensis IDA1967 (NR_029002.1)	99.65	$30 \pm 1$	$100 \pm 0$	$100 \pm 0$	7 ± 1	56 ± 10	69 ± 6
10	Bacillus aryabhattai	<i>B. aryabhattai</i> B8W22 (NR_115953.1)	99.87	89 ± 11	$100 \pm 0$	$100 \pm 0$	4 ± 1	24 ± 6	90 ± 10
11	Bacillus aryabhattai	<i>B. aryabhattai</i> B8W22 (NR_115953.1)	99.80	$24 \pm 5$	$100 \pm 0$	$100 \pm 0$	$10 \pm 1$	26 ± 4	81 ± 11
12	Bacillus oryzisoli	<i>B. oryzisoli</i> 1DS3-10 (NR_151979.1)	98.88	89 ± 2	98 ± 2	97 ± 3	7 ± 1	8 ± 2	26 ± 5
13	Bacillus cereus	B. cereus CCM 2010 (NR_115714.1)	99.74	$100 \pm 0$	100 ± 0	100 ± 0	7 ± 4	42 ± 3	92 ± 8
14	Bacillus cereus	B. cereus CCM 2010 (NR_115714.1)	99.01	$100 \pm 0$	100 ± 0	100 ± 0	47 ± 5	75 ± 4	75 ± 4
15	Bacillus chungangensis	B. chungangensis CAU 348 (NR_116709.1)	99.32	15 ± 5	93 ± 8	100 ± 0	9 ± 2	9 ± 2	22 ± 1
16	Bacillus cohnii	<i>B. cohnii</i> DSM 6307 (NR_026138.1)	99.67	48 ± 5	100 ± 0	100 ± 0	11 ± 1	13 ± 2	$35 \pm 4$
17	Bacillus cohnii	<i>B. cohnii</i> DSM 6307 (NR_026138.1)	99.40	9 ± 3	41 ± 2	85 ± 4	9 ± 1	10 ± 1	14 ± 1
18	Bacillus cohnii	<i>B. cohnii</i> DSM 6307 (NR_026138.1)	99.60	21 ± 3	100 ± 0	100 ± 0	8 ± 2	13 ± 2	37 ± 7
19	Bacillus cohnii	<i>B. cohnii</i> DSM 6307 (NR_026138.1)	99.67	57 ± 12	100 ± 0	100 ± 0	9±3	18 ± 6	81 ± 11
20	Bacillus firmus	B. firmus IAM 12464 (NR_112635.1)	99.13	96 ± 4	100 ± 0	100 ± 0	8 ± 1	10 ± 2	24 ± 3
21	Bacillus depressus	B. depressus BZ1 (NR_146034.1)	99.52	50 ± 6	100 ± 0	100 ± 0	9 ± 1	26 ± 1	100 ± 0
22	Bacillus depressus	B. depressus BZ1 (NR_146034.1)	99.59	18 ± 5	100 ± 0	100 ± 0	7 ± 3	13 ± 1	21 ± 6
23	Bacillus halodurans	B. halodurans DSM 497 (NR_025446.1)	99.87	18 ± 5	100 ± 0	100 ± 0	7 ± 1	11 ± 2	21 ± 1
24	Bacillus horikoshii	B. horikoshii DSM 8719 (NR_040852.1)	99.27	25 ± 5	100 ± 0	100 ± 0	8 ± 0	11 ± 1	53 ± 7
25	Bacillus horneckiae	B. horneckiae 1P01SC (NR_116474.1)	99.34	33 ± 5	100 ± 0	100 ± 0	9 ± 4	11 ± 5	25 ± 11
26	Bacillus horneckiae	B. horneckiae 1P01SC (NR_116474.1)	99.78	40 ± 5	100 ± 0	100 ± 0	3 ± 2	11 ± 1	34 ± 3
27	Bacillus licheniformis	B. licheniformis DSM 13 (NR_118996.1)	99.54	41 ± 4	100 ± 0	100 ± 0	8±6	22 ± 4	51 ± 7
28	Bacillus licheniformis	B. licheniformis DSM 13 (NR_118996.1)	99.54	77 ± 8	99 ± 1	100 ± 0	12 ± 3	20 ± 3	59 ± 4
29	Bacillus litoralis	B. litoralis SW-211 (NR_043015.1)	98.60	44 ± 12	100 ± 0	100 ± 0	12 ± 4	18 ± 8	96 ± 4
30	Bacillus mannanilyticus	B. mannanilyticus AM-001 (NR_040851.1)	96.93	30 ± 6	100 ± 0	100 ± 0	10 ± 0	11 ± 1	25 ± 6
31	Bacillus megaterium	<i>B. megaterium</i> NBRC 15308 (NR_112636.1)	99.86	100 ± 0	100 ± 0	100 ± 0	7 ± 2	22 ± 2	92 ± 8
32	Bacillus niacini	B. niacini IFO15566 (NR_024695.1)	98.94	24 ± 10	100 ± 0	100 ± 0	8 ± 1	50 ± 10	98 ± 1
33	Bacillus bataviensis	B. bataviensis IDA1115 (NR_036766.1)	97.32	95 ± 5	100 ± 0	100 ± 0	9±3	59 ± 5	98 ± 2
34	Bacillus acidiceler	B. acidiceler CBD 119 (NR_043774.1)	99.67	42 ± 7	100 ± 0	100 ± 0	7 ± 2	9 ± 2	15 ± 2
35	Bacillus australimaris	<i>B. australimaris</i> MCCC 1A05787 (NR_148787.1)	99.67	100 ± 0	100 ± 0	100 ± 0	13 ± 3	10 ± 1	48 ± 3
36	Bacillus xiamenensis	<i>B. xiamenensis</i> MCCC 1A00008 (NR_148244.1)	99.47	100 ± 0	100 ± 0	100 ± 0	7 ± 1	12 ± 2	48 ± 1
37	Bacillus xiamenensis	<i>B. xiamenensis</i> MCCC 1A00008 (NR_148244.1)	98.81	100 ± 0	100 ± 0	100 ± 0	10 ± 4	12 ± 1	62 ± 1
38	Bacillus xiamenensis	<i>B. xiamenensis</i> MCCC 1A00008 (NR_ 148244.1)	99.74	96 ± 4	100 ± 0	100 ± 0	10 ± 1	16 ± 2	63 ± 2

# Table 1 Degradation (%) of Methyl Red and Orange II by 79 bacterial isolates<sup>a</sup> (Continued)

Isolate	Identity based on 16S rRNA	BLAST on NCBI 16S RefSeq database	2	Methyl	Red <sup>b</sup>		Orange	e II <sup>b</sup>	
#		Type strain (Sequence ID)	% Identity	1 h	8 h	24 h	1 h	8 h	24 h
39	Bacillus simplex	B. simplex NBRC 15720 (NR_042136.1)	99.34	72 ± 8	$100 \pm 0$	$100 \pm 0$	18 ± 2	74 ± 7	74 ± 7
40	Bacillus subtilis	B. subtilis IAM 12118 (NR_112116.2)	99.60	97 ± 3	$100 \pm 0$	$100 \pm 0$	$11 \pm 0$	$11 \pm 3$	$20 \pm 3$
41	Bacillus subtilis	B. subtilis IAM 12118 (NR_112116.2)	99.47	89 ± 5	$100 \pm 0$	$100 \pm 0$	8 ± 1	$14 \pm 1$	$45 \pm 3$
42	Bacillus subtilis	B. subtilis IAM 12118 (NR_112116.2)	99.34	$100 \pm 0$	$100 \pm 0$	$100 \pm 0$	7 ± 2	8 ± 1	19±3
43	Bacillus thermoamylovorans	<i>B. thermoamylovorans</i> LMG 18084 (NR_117028.1)	99.60	19 ± 5	98 ± 2	99 ± 1	13 ± 3	97 ± 3	98 ± 1
44	Bacillus thuringiensis	B. thuringiensis IAM 12077 (NR_043403.1)	99.87	$100 \pm 0$	$100 \pm 0$	$100 \pm 0$	$12 \pm 0$	$56 \pm 3$	95 ± 2
45	Bacillus thuringiensis	B. cereus CCM 2010 ( NR_115714.1)	99.80	96 ± 4	$100 \pm 0$	$100 \pm 0$	4 ± 2	44 ± 9	91 ± 5
46	Barrientosiimonas humi	B. humi 39 (NR_126227.1)	99.80	13 ± 1	18 ± 1	$33 \pm 2$	9 ± 2	9 ± 2	11 ± 2
47	Brachybacterium faecium	<i>B. faecium</i> DSM 4810 (NR_074655.2)	97.78	14 ± 2	$100 \pm 0$	$100 \pm 0$	11 ± 1	$13 \pm 3$	76 ± 12
48	Brachybacterium paraconglomeratum	<i>B. paraconglomeratum</i> LMG 19861 (NR_025502.1)	99.12	14 ± 5	44 ± 8	90 ± 7	7 ± 2	10 ± 2	16 ± 5
49	Brevibacillus formosus	B. formosus DSM 9885 (NR_040979.1)	99.32	36 ± 4	$100 \pm 0$	$100 \pm 0$	$10 \pm 2$	$13 \pm 1$	$33 \pm 2$
50	Brevibacillus laterosporus	B. laterosporus DSM 25 (NR_112212.1)	99.53	82 ± 8	77 ± 0	$100 \pm 0$	9 ± 1	99 ± 1	$100 \pm 0$
51	Brevibacillus parabrevis	<i>B. parabrevis</i> IFO 12334 (NR_040981.1)	99.26	$28 \pm 10$	83 ± 6	100 ± 0	9 ± 1	14 ± 2	71 ± 3
52	Dermacoccus barathri	D. barathri MT2.1 (NR_043261.1)	99.79	$14 \pm 1$	46 ± 4	99 ± 1	9 ± 2	11 ± 1	13 ± 2
53	Pseudomonas aeruginosa	P. aeruginosa DSM 50071 (NR_117678.1)	99.93	20 ± 5	91 ± 10	99 ± 1	10 ± 5	21 ± 10	94 ± 6
54	Fictibacillus arsenicus	F. phosphorivorans Ca7 (NR_118455)	99.38	$24 \pm 5$	30 ± 3	40 ± 3	13 ± 3	16 ± 2	36 ± 10
55	Fictibacillus phosphorivorans	F. phosphorivorans Ca7 (NR_118455)	99.58	31 ± 9	94 ± 6	96 ± 3	9 ± 1	10 ± 3	29 ± 9
56	Kocuria marina	K. marina KMM 3905 (NR_025723.1)	99.86	9 ± 4	54 ± 8	100 ± 0	7 ± 4	9 ± 2	11 ± 2
57	Lysinibacillus fusiformis	L. fusiformis DSM 2898 (NR_042072.1)	99.93	32 ± 8	100 ± 0	100 ± 0	9±3	11 ± 3	83 ± 4
58	Lysinibacillus fusiformis	L. fusiformis DSM 2898 (NR_042072.1)	99.80	74 ± 11	100 ± 0	100 ± 0	4 ± 1	10 ± 3	82 ± 8
59	Lysinibacillus fusiformis	L. fusiformis DSM 2898 (NR_042072.1)	99.32	29 ± 10	$100 \pm 0$	100 ± 0	8 ± 3	12 ± 2	83 ± 4
60	Lysinibacillus macroides	L. macroides LMG 18474 (NR_114920.1)	99.40	44 ± 7	100 ± 0	100 ± 0	8 ± 3	12 ± 3	80 ± 3
61	Lysinibacillus macroides	L. macroides LMG 18474 (NR_114920.1)	99.33	38 ± 8	$100 \pm 0$	100 ± 0	5 ± 2	8 ± 3	85 ± 2
62	Micrococcus luteus	M. luteus NCTC 2665 ( NR_075062.2)	99.53	8 ± 2	42 ± 7	97 ± 3	6 ± 1	11 ± 3	18 ± 3
63	Micrococcus luteus	M. luteus NCTC 2665 ( NR_075062.2)	98.05	10 ± 2	34 ± 5	89 ± 5	7 ± 1	9 ± 2	17 ± 2
64	Oceanobacillus chironomi	O. chironomi T3944D ( NR_043700.1)	97.14	$24 \pm 6$	$100 \pm 0$	100 ± 0	13 ± 4	87 ± 9	92 ± 8
65	Paenibacillus amylolyticus	P. amylolyticus NRRL NRS-290 (NR_025882.1)	99.27	94 ± 6	100 ± 0	100 ± 0	1 ± 0	17 ± 6	60 ± 2
66	Paenibacillus tundrae	P. tundrae A10b ( NR_044525.1)	99.27	$100 \pm 0$	$100 \pm 0$	$100 \pm 0$	4 ± 1	$15 \pm 4$	53 ± 1
67	Paenibacillus tundrae	P. tundrae A10b ( NR_044525.1)	99.67	89 ± 9	$100 \pm 0$	$100 \pm 0$	6 ± 2	12 ± 2	51 ± 9
68	Paenibacillus glucanolyticus	P. glucanolyticus DSM 5162 (NR_ 040883.1)	99.47	16 ± 1	100 ± 0	100 ± 0	8 ± 2	10 ± 1	46 ± 3
69	Paenibacillus lautus	P. lautus JCM 9073 (NR_040882.1)	99.47	96 ± 4	$100 \pm 0$	$100 \pm 0$	6 ± 4	$100 \pm 0$	$100 \pm 0$
70	Paenibacillus lautus	P. lautus JCM 9073 (NR_040882.1)	99.60	19 ± 3	96 ± 4	97 ± 4	10 ± 1	$100 \pm 0$	$100 \pm 0$
71	Providencia vermicola	P. vermicola OP1 ( NR_042415.1)	99.66	44 ± 10	$100 \pm 0$	$100 \pm 0$	7 ± 4	9 ± 2	22 ± 2
72	Pseudomonas aeruginosa	P. aeruginosa DSM 50071 (NR_117678.1)	99.73	11 ± 2	29 ± 2	100 ± 0	11 ± 2	12 ± 2	97 ± 2
73	Pseudomonas aeruginosa	P. aeruginosa DSM 50071 ( NR_117678.1)	99.87	10 ± 3	41 ± 3	84 ± 7	7 ± 1	11 ± 0	16 ± 5
74	Pseudomonas aeruginosa	P. aeruginosa DSM 50071 (NR_117678.1)	99.93	7 ± 2	$28 \pm 6$	$100 \pm 0$	7 ± 1	$14 \pm 4$	100 ± 0

### Table 1 Degradation (%) of Methyl Red and Orange II by 79 bacterial isolates<sup>a</sup> (Continued)

Isolate	Identity based on 16S rRNA	BLAST on NCBI 16S RefSeq database	2	Methyl	Red <sup>b</sup>		Orange	e II <sup>b</sup>	
#		Type strain (Sequence ID)	% Identity	1 h	8 h	24 h	1 h	8 h	24 h
75	Pseudomonas aeruginosa	P. aeruginosa DSM 50071 (NR_117678.1)	99.87	9 ± 3	31 ± 9	$100 \pm 0$	11 ± 1	18 ± 5	100 ± 0
76	Lysinibacillus macroides	L macroides LMG 18474 (NR_114920.1)	99.33	51 ± 3	$100 \pm 0$	100 ± 0	7 ± 4	13 ± 1	89 ± 3
77	Pseudomonas aeruginosa	P. aeruginosa DSM 50071 (NR_117678.1)	99.87	6 ± 1	$28 \pm 6$	100 ± 0	4 ± 1	15 ± 3	$100 \pm 0$
78	Psychrobacillus lasiicapitis	P. lasiicapitis NEAU-3TGS17 (NR_159144.1)	98.95	16 ± 2	92 ± 8	100 ± 0	5 ± 2	13 ± 2	82 ± 13
79	Roseomonas mucosa	R. mucosa MDA5527 ( NR_028857.1)	99.72	13 ± 3	38 ± 5	97 ± 2	10 ± 2	9 ± 2	17 ± 3
80	Sporosarcina luteola	<i>S. luteola</i> Y1 ( NR_112844.1)	99.67	11 ± 2	71 ± 7	100 ± 0	5 ± 2	11 ± 3	70 ± 14
81	Sporosarcina soli	S. soli 180 ( NR_043527.1)	99.66	9 ± 1	90 ± 7	100 ± 0	12 ± 1	12 ± 2	27 ± 8

<sup>a</sup>The data show the mean of degradation (%)  $\pm$  SD from repeated triplicated assay using 96-well microphates

<sup>b</sup>100 μM Orange II and 100 μM Methyl Red were used in the degradation assay, except that 50 μM and 25 μM Methyl Red were used for isolates #4 and #16, respectively and 12.5 μM Methyl Red was used for isolates #12 and #20. Absorbance wavelengths for Methyl Red and Orange II were 430 nm and 475 nm, respectively

experiment was triplicated and repeated twice. The final degradation (%) is the mean of data from repeated triplicated assay.

# Multiple sequence alignment, construction of a phylogenetic tree, and tree annotation

The multiple sequence alignment and construction of a neighbor-joining tree of 16S rDNA sequences from 79 bacterial isolates were performed using Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/), a new multiple sequence alignment program, that uses seeded guide trees and HMM profile-profile techniques to generate alignments between sequences, with the default settings. To annotate the neighbor-joining tree with degradation rate (%) of Methyl Red and Orange II by 79 bacterial isolates (i.e., Fig. 2, a tree with multi-value bar chart), iTOL (https://itol.embl.de/) was used: input datasets of a neighbor-joining tree in Newick format and a dataset\_multibar\_template.txt updated with the degradation data of Methyl Red and Orange II by 79 bacterial isolates (Table 1). A template txt file (dataset\_multibar\_ template.txt) can be downloaded from the iTOL website (https://itol.embl.de/help.cgi#datasets).

## **Results and discussion**

A two-step strategy was used in this study; initially degradation of azo dye was tested for all tattoo inkderived bacterial isolates using Methyl Red and Orange II (step 1), followed by an evaluation of selected bacteria with respect to the degradation of 11 representative azo dyes, including 3 commercial tattoo ink azo dyes (step 2) (Fig. 1).

# Step 1: Screen 79 bacterial isolates for azo dye degradability using Methyl Red and Orange II

A total of 79 bacterial isolates were tested for the decolorization of azo dyes (Fig. 2 and Table 1). Methyl Red and Orange II were used as model azo dye compounds based on their molecular weight. Overall, results of the screening assay showed that most of these tattoo

Table 2 Degradation (%) of 8 structurally diverse azo dyes within 48 h by selected bacterial isolates

Azo dyes	MW	$\mu^{a}$	nm <sup>b</sup>	Azo dye de	gradatio	n (%) in bact	erial strains (isolate #) <sup>c</sup>						
				B. depressus (#21)	B. niacini (#32)	B. bataviensis (#33)	B. thermosamylovorans (#43)	Brevi. laterosporus (#50)	Paeni. lautus (#69)	Paeni. lautus (#70)	P. aeruginosa (#74)	P. aeruginosa (#75)	P. aeruginosa (#77)
Sudan I	248.3	100	475	86 ± 5	88 ± 2	84 ± 3	85 ± 1	87 ± 1	87 ± 5	87 ± 1	88 ± 5	87 ± 7	90 ± 2
Sudan II	276.3	75	493	39 ± 2	38 ± 1	28 ± 2	40 ± 1	39 ± 6	37 ± 1	39 ± 0	42± 1	40 ± 3	46 ± 0
Sudan III	352.4	45	510	77 ± 1	67 ± 1	71 ± 2	51 ± 2	55 ± 7	85 ± 7	71 ± 9	34 ± 1	32 ± 8	29 ± 8
Sudan IV	380.5	45	520	25 ± 1	32 ± 3	39 ± 9	32 ± 0	26 ± 4	$40 \pm 6$	34 ± 6	20 ± 7	21 ± 4	32 ± 1
Orange G	452.4	100	475	16 ± 7	6 ± 1	26 ± 6	97 ± 1	94 ± 2	97 ± 3	97 ± 1	36 ± 4	35 ± 5	36 ± 2
Amaranth	452.4	45	520	59 ± 8	72 ± 1	59 ± 9	99 ± 1	99 ± 1	99 ± 2	99 ± 0	70 ± 1	85 ± 1	89 ± 8
Ponceau BS	556.5	100	510	94 ± 1	92 ± 1	90 ± 6	94 ± 4	92 ± 1	96 ± 2	97 ± 1	97 ± 1	96 ± 1	96 ± 1
Direct Blue 15	992.8	45	620	23 ± 2	23 ± 6	21 ± 5	93 ± 0	86 ± 4	96 ± 0	97 ± 1	39 ± 5	37 ± 7	35 ± 3

<sup>a</sup>Concentration of azo dyes

<sup>b</sup>Absorbance wavelength measured for respective azo dyes

<sup>c</sup>The data show the mean of degradation (%)  $\pm$  SD from repeated triplicated assay using 15-mL concical centrifuge tubes

and PMU ink-derived bacteria well degrade (or decolorize) the low molecular weight azo dye, Methyl Red. Sixty two (78%) and 74 (93%) out of 79 isolates degraded more than 90% of Methyl Red at 8 and 24 h after incubation, respectively (Fig. 2 and Table 1). The remaining five isolates showed degradation of Methyl Red, ranged from 33 to 89% after 24 h incubation. Although there were no previous high-throughput screening data to compare, the results showed that all of the tested bacterial isolates have an ability of azo dye degradation. It indicates that there might be a functional correlation of the capability of azo dye degradation of



these bacteria with tattoo inks, an azo dye-rich environment. In case of the high molecular weight azo dye, Orange II, 20 isolates (25%) degraded the dye with > 90% after 24 h incubation while 4 isolates achieved 90% of degradation at 4 h post-incubation (Table 1). The results showed that degradation of Methyl Red was much faster than Orange II, which indicates that the low molecular weight azo dye may be degraded by microorganisms quickly compared with the high molecular weight azo dye (Table 1).

Taxonomically, a wide range of bacterial species were previously reported for their ability to degrade azo dyes, which included bacterial genera Micrococcus, Geobacillus, Pseudomonas, Kocuria, Sphingomonas, Bacillus, Shewanella, Acinetobacter, Staphylococcus, Aeromonas, Corynebacterium, Dermacoccus, and Kocuria, to name a few (Saratale et al. 2011; Solís et al. 2012; Stingley et al. 2010). However, in this study, we additionally identified the ability of azo dye degradation in the bacterial genera, Brachybacterium, Desemzia, Psychrobacillus, Fictibacillus, Roseomonas, and Sporosarcina, which have not been previously reported to degrade azo dyes. Generally, isolates which belonged to the same taxonomy in most cases had abilities of azo dye degradation at a comparable level. However, we observed that 4 strains of B. cohnii degraded Orange II at different levels, ranged from 14 to 81%. Strains of B. depressus and P. aeruginosa also showed a significant difference in the level of Orange II degradation (Table 1). It indicated that degradation capability of certain azo dyes depends on the characteristics of each individual strains regardless of its taxonomy.

# Step 2: Evaluation of the selected bacterial isolates for their degradation capability for 11 representative azo dyes, including 3 commercial tattoo ink azo dyes

In step 2, based on the activity of Methyl Red and Orange II degradation in step 1, we selected 10 bacterial isolates. We then further evaluated their ability of azo dye degradation in detail using a broad range of azo dyes (Fig. 1 and Tables 2 and 3). A representative of 11 structurally diverse azo dyes, including 3 commercial tattoo ink azo dyes, was selected for this purpose (Fig. 1). It included, in the order of increasing molecular weight, Sudan I, Solvent Red 1, Sudan II, Sudan III, Sudan IV, Lithol Rubin BK, Orange G, Amaranth, Alphamine Red R, Ponceau BS, and Direct Blue 15. They were selected based on molecular weight (MW 248.3 to 992.8), number of azo bond (1-2 azo bonds), number of benzene ring (2–6 benzene rings), presence of substituted (sulfonated) benzene ring, and the usage as tattoo ink pigments.

The 10 tested bacterial strains belonged to *Bacillus* species ( $\times$  4), *Brevi. laterosporus* ( $\times$  1), *Paeni. lautus* ( $\times$  2), and *P. aeruginosa* ( $\times$  3). Overall, all 10 isolates had a

varying ability to degrade the 8 structurally diverse representative azo dyes (Fig. 1 and Table 2). The 10 isolates showed a relatively high degradation rate of Sudan I, Amaranth, and Ponceau BS (> 59%) but not for Sudan II and Sudan IV (< 40%). On the other hand, Orange G, Direct Blue 15, and Sudan III were well degraded by some isolates but not well degraded by other isolates. *Paeni. lautus, Brevi. laterosporus,* and *B. thermoamylovorans* were able to degrade more than 86% of Orange G and Direct Blue 15. On the other hand, only isolates, belonging to *Bacillus* and *Bacillus*-related genera, showed a relatively high degradation (51–85%) toward Sudan III.

Different bacteria have been reported to decolorize azo dyes tested in this study. Skin bacteria belonging to the genera Staphylococcus, Corynebacterium, Micrococcus, Dermacoccus, and Kocuria, were able to decolorize Methyl Red in nutrient broth within 24 h with 74-100% reduction under static condition (Stingley et al. 2010). In our study, we had strains belonging to Dermacoccus and Kocuria genera, which had achieved over 99% reduction of Methyl Red in TSB by 24 h (Table 1). A strains of Brevi. laterosporus, was found to decolorize 93% of Methyl Red within 12 h in nutrient broth (Gomare and Govindwar 2009). Complete decolorization of Mtehyl Red was observed in our study within 24 h in TSB under static condition by a strain of Brevi. laterosporus (Table 1). Among those azo dyes tested in step 2, only one case exists in the literature where a *P. aeruginosa* strain was able to decolorize 96% of Amaranth in distilled water within 6 h (Jadhav et al. 2013). Whereas, in our study, 3 different strains of P. aeruginosa were capable of decolorizing Amaranth in TSB with the range of reduction rate between 70 and 89% within 48 h (Table 2).

In case of azo dyes widely used as pigments in the commercial tattoo ink products, the 10 isolates degraded more than 74% of Lithol Rubin BK and Alphamine Red R (Fig. 1 and Table 3). On the other hand, only *Brevi. Laterosporus* (#50) showed a high degradation rate (95%) to Solvent Red 1; other isolates showed a degradation rate around or less than 50% (Fig. 1 and Table 3). There are no reports available on the decolorization of azo dyes used in tattoo inks. Thus, the data obtained in this study may contribute to decolorization of tattoo ink azo dyes by pure bacterial cultures.

Overall, the results of this study suggest that there is a general dependency on azo dye structure and bacterial taxonomy and pleiotropic and epistatic functional interactions among diverse azo dye degrading enzymes in the degradation of azo dyes. Out of the tested 10 isolates, *B. thermoamylovorans* (isolate #43), *Brevi. laterosporus* (#50), and 2 strains of *Paeni. lautus* (#69 and #70) were the top 4 bacteria with high efficiencies of azo dye degradation. High degradation ability of azo dyes by

				Azo dye degra	dation (%) in	bacterial strai	ns (isolate #) <sup>c</sup>						
Azo dyes	MM	е Ми	a mu	B. depressus (#21)	B. niacini (#32)	B. bataviensis (#33`)	B. thermosamylovorans (#43)	Brevi. laterosporus (#50)	Paeni. lautus (#69)	Paeni. lautus (#70)	P. aeruginosa (#74)	P. aeruginosa (#75)	P. aeruginosa (#77)
Solvent Red 1	278.3	100	500	46 ± 9	58 ± 9	47 ± 13	53 ± 8	95 ± 6	53 ± 15	58 ± 18	47 ± 18	45 ± 15	39 ± 15
Lithol Rubin BK	430.3	100	500	99 ± 1	99 ± 1	97 ± 2	81 ± 19	96 ± 3	98 ± 2	100 ± 0	79 ± 9	76 ± 5	74 ± 9
Alphamine Red R	533.6	100	475	92 ± 9 <sup>d</sup>	97 ± 1	98 ± 2	94 ± 0	99 ± 1 <sup>d</sup>	$100 \pm 0^{dd}$	98 ± 4 <sup>d</sup>	89 ± 2	88 ± 2	88 ± 6
<sup>a</sup> Concentration c	of azo d	lyes											

Table 3 Degradation (%) of 3 azo dyes used widely as pigments in the commercial tattoo ink products by selected bacterial isolates

Concentration of all upper babsorbenetions are used for respective all dynamics of the babsorbenetic measured for respective all distances <sup>c</sup>The data show the mean of degradation (%)  $\pm$  SD from repeated triplicated assay using 96-well microphates <sup>d</sup>Tested with 50 µM of Alphamine Red R

members of the genera *Brevibacillus* and *Paenibacillus* have been shown (Alhassani et al. 2007; Gomare and Govindwar 2009; Nawahwi et al. 2013; Ramya et al. 2008; Srinivasan et al. 2014).

## Conclusions

This study was a novel attempt to (i) examine the capability of azo dye degradation (or decolorization) of bacterial strains isolated from tattoo inks, which provide an azo dye-rich environment, and (ii) introduce a highthroughput azo dye degradation assay to identify azo dye-degrading bacteria. The study confirmed the ability of phylogenetically diverse bacteria, isolated from azo dye-rich tattoo inks, to degrade a diverse range of azo dyes, including 3 azo dyes used in commercial tattoo inks. Four bacterial isolates, belonging to B. thermoamylovorans (#43), Brevi. laterosporus (#50), and Paeni. lautus (#69 and #70), exhibited an excellent capability of degradation of a diverse range of azo dyes. Further studies, including genome sequencing and functional genomics with these isolates, should be followed for a systematic understanding of the mechanism of azo dye degradation in these microorganisms, which is essential for their practical use in bioremediation application for removal of azo dyes via comparative insights on the differences in properties from the azo pigment-degrading bacteria that have been reported so far.

#### Acknowledgements

We thank Nakissa Sadrieh, Steven L. Foley, and Bruce D. Erickson for critical review of the manuscript. This work was supported in part by an appointment to the Postgraduate Research Fellowship Program at the National Center for Toxicological Research, administered by the Oak Ridge Institute for Science and Education through an interagency agreement between the U. S. Department of Energy and the U. S. Food and Drug Administration. Authors declare that there is no conflicts of interest. The opinions and views presented in this article do not necessarily reflect current or future opinions or policies of the U.S. FDA.

#### Authors' contributions

SWN, XWC, JJ, and OK performed the experiments. OK and SJK designed the study, analyzed the data, and wrote the manuscript. HC made contributions to the analysis of experimental data and revised the manuscript. MSM helped in analyzing the data and revised the manuscript. SK and CEC supervised the project. All authors read and approved the final manuscript.

#### Funding

Not applicable

#### Availability of data and materials

Data is available under request.

#### Declarations

Ethics approval and consent to participate Not applicable

### Consent for publication

Not applicable.

### **Competing interests**

The authors declare that they have no competing interests.

#### Author details

<sup>1</sup>Division of Microbiology, National Center for Toxicological Research, U.S. Food and Drug Administration, Jefferson, AR, USA. <sup>2</sup>Present address: Sichuan Institute for Food and Drug Control, Chengdu 611731, Sichuan, China. <sup>3</sup>Office of Cosmetics and Colors, Center for Food Safety and Applied Nutrition, U.S. Food and Drug Administration, College Park, MD, USA.

#### Received: 18 May 2021 Accepted: 21 August 2021 Published online: 21 September 2021

#### References

- Alhassani HA, Rauf MA, Ashraf SS (2007) Efficient microbial degradation of toluidine blue dye by *Brevibacillus* sp. Dyes Pigments 75(2):395–400. https:// doi.org/10.1016/j.dyepig.2006.06.019
- Bäumler W, Eibler ET, Hohenleutner U, Sens B, Sauer J, Landthaler M (2000) Qswitch laser and tattoo pigments: first results of the chemical and photophysical analysis of 41 compounds. Lasers Surg Med 26(1):13–21. https://doi.org/10.1002/(SICI)1096-9101(2000)26:1<13::AID-LSM4>3.0.CO;2-S
- Carmen Z, Daniela S (2012) Textile organic dyes-characteristics, polluting effects and separation/elimination procedures from industrial effulents - a critical overview. In: Puzyn T (ed) Organic Pollutants Ten Years after the Sockholm Convention - Environmental and Analytical Update,, pp 55–86
- Chung KT (2016) Azo dyes and human health: a review. J Environ Sci Health C Environ Carcinog Ecotoxicol Rev 34:233–261
- Feng J, Cerniglia CE, Chen H (2012) Toxicological significance of azo dye metabolism by human intestinal microbiota. Front Biosci 4:568–586
- Forgacs E, Cserhati T, Oros G (2004) Removal of synthetic dyes from wastewaters: a review. Environ Int 30(7):953–971. https://doi.org/10.1016/j.envint.2004.02. 001
- Gomare SS, Govindwar SP (2009) *Brevibacillus laterosporus* MTCC 2298: a potential azo dye degrader. J Appl Microbiol 106:993-1004, 3, DOI: https://doi.org/1 0.1111/j.1365-2672.2008.04066.x
- Jadhav SB, Patil NS, Watharkar AD, Apine OA, Jadhav JP (2013) Batch and continuous biodegradation of Amaranth in plain distilled water by P. aeruginosa BCH and toxicological scrutiny using oxidative stress studies. Environ Sci Pollut Res Int 20(5):2854–2866. https://doi.org/10.1007/s11356-012-1155-y
- Jin XC, Liu GQ, Xu ZH, Tao WY (2007) Decolorization of a dye industry effluent by Aspergillus fumigatus XC6. Appl Microbiol Biotechnol 74(1):239–243. https:// doi.org/10.1007/s00253-006-0658-1
- Meehan C, Bjourson AJ, McMullan G (2001) Paenibacillus azoreducens sp. nov., a synthetic azo dye decolorizing bacterium from industrial wastewater. Int J Syst Evol Microbiol 51:1681-1685, 5, DOI: https://doi.org/10.1099/00207713-51-5-1681
- Nawahwi MZ, Ibrahim Z, Yahya A (2013) Degradation of the azo dye reactive red 195 by *Paenibacillus* spp. R2. J Bioremed Biodeg 4:174
- Nho SW, Kim M, Kweon O, Kim SJ, Moon MS, Periz G, Huang MJ, Dewan K, Sadrieh NK, Cerniglia CE (2020) Microbial contamination of tattoo and permanent makeup inks marketed in the US: a follow-up study. Lett Appl Microbiol 71(4):351–358. https://doi.org/10.1111/lam.13353
- Nho SW, Kim SJ, Kweon O, Howard PC, Moon MS, Sadrieh NK, Cerniglia CE (2018) Microbiological survey of commercial tattoo and permanent makeup inks available in the United States. J Appl Microbiol 124(5):1294–1302. https://doi. org/10.1111/jam.13713
- Ramya M, Anusha B, Kalavathy S (2008) Decolorization and biodegradation of Indigo carmine by a textile soil isolate *Paenibacillus larvae*. Biodegradation 19(2):283–291. https://doi.org/10.1007/s10532-007-9134-6
- Rawat D, Mishra V, Sharma RS (2016) Detoxification of azo dyes in the context of environmental processes. Chemosphere 155:591–605. https://doi.org/10.101 6/j.chemosphere.2016.04.068
- Robinson T, McMullan G, Marchant R, Nigam P (2001) Remediation of dyes in textile effluent: a critical review on current treatment technologies with a proposed alternative. Bioresour Technol 77(3):247–255. https://doi.org/10.101 6/S0960-8524(00)00080-8
- Saratale RG, Saratale GD, Chang JS, Govindwar SP (2011) Bacterial decolorization and degradation of azo dyes: A review. J Taiwan Instit Chem Eng 42(1):138– 157. https://doi.org/10.1016/j.jtice.2010.06.006
- Solís M, Solís A, Pérez HI, Manjarrez N, Flores M (2012) Microbial decolouration of azo dyes: A review. Process Biochem 47(12):1723–1748. https://doi.org/10.101 6/j.procbio.2012.08.014

- Srinivasan GP, Sikkanthar A, Elamaran A, Delma CR, Subramaniyan K, Somasundaram ST (2014) Biodegradation of carcinogenic textile azo dyes using bacterial isolates of mangrove sediment. J Costal Life Med 2:154–162
- Stingley RL, Zou W, Heinze TM, Chen H, Cerniglia CE (2010) Metabolism of azo dyes by human skin microbiota. J Med Microbiol 59(1):108–114. https://doi. org/10.1099/jmm.0.012617-0
- Stolz A (2001) Basic and applied aspects in the microbial degradation of azo dyes. Appl Microbiol Biotechnol 56(1-2):69–80. https://doi.org/10.1007/s00253 0100686
- Telke AA, Joshi SM, Jadhav SU, Tamboli DP, Govindwar SP (2010) Decolorization and detoxification of Congo red and textile industry effluent by an isolated bacterium *Pseudomonas* sp. SU-EBT Biodegradation 21(2):283–296. https:// doi.org/10.1007/s10532-009-9300-0
- Xu M, Guo J, Cen Y, Zhong X, Cao W, Sun G (2005) Shewanella decolorationis sp. nov., a dye-decolorizing bacterium isolated from activated sludge of a wastewater treatment plant. Int J Syst Evol Microbiol 55:363-368, 1, DOI: https://doi. org/10.1099/ijs.0.63157-0

### **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

#### Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

#### At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

